



# CHEMICAL AND FOOD PROCESSES APPLIED TO GOAT'S MILK: NON-CONVENTIONAL DERIVATIVES

Eduardo Zorrilla Tarazona - Juan Luis Pérez  
Marín - Ingrid Yossy Robles Castañeda - Ader  
Martín Guerra Rios - Raúl Tello Suarez - José  
Antonio Luna García - Carlos Ruíz Padilla

URUGUAY, 2025

ISBN: 978-9915-9752-9-0



9 789915 975290

# Chemical and food processes applied to goat's milk: Non-Conventional Derivatives

Eduardo Zorrilla Tarazona, Juan Luis Pérez Marín, Ingrid Yossy Robles Castañeda, Ader Martín Guerra Rios, Raúl Tello Suarez, José Antonio Luna García, Carlos Ruíz Padilla

© Eduardo Zorrilla Tarazona, Juan Luis Pérez Marín, Ingrid Yossy Robles Castañeda, Ader Martín Guerra Rios, Raúl Tello Suarez, José Antonio Luna García, Carlos Ruíz Padilla, 2025

First edition: February, 2025

Edited by:

Editorial Mar Caribe

[www.editorialmarcaribe.es](http://www.editorialmarcaribe.es)

Av. General Flores 547, Colonia, Colonia-Uruguay.

Cover Design: Yelitza Sánchez Cáceres

E-book available at: <https://editorialmarcaribe.es/ark:/10951/isbn.9789915975290>

Format: electronic

ISBN: 978-9915-9752-9-0

ARK: [ark:/10951/isbn.9789915975290](https://editorialmarcaribe.es/ark:/10951/isbn.9789915975290)

## Non-Commercial Attribution Rights Notice:

Authors may authorize the general public to reuse their works solely for non-profit purposes, readers may use one work to generate another work, as long as research credit is given, and they grant the publisher the right to first publish their essay under the terms of the [CC BY-NC 4.0 license](https://creativecommons.org/licenses/by-nc/4.0/).

## Editorial Mar Caribe, signatory No. 795 of 12.08.2024 of the Declaration of Berlin:

*"... We feel compelled to address the challenges of the internet as an emerging functional medium for the distribution of knowledge. Obviously, these advances may significantly modify the nature of scientific publishing, as well as the existing system of quality assurance..."* (Max Planck Society, ed. 2003., pp. 152-153).

## [Editorial Mar Caribe-Member of OASPA:](#)

As a member of the Open Access Scholarly Publishing Association, we support open access in accordance with [OASPA's](#) code of conduct, transparency, and best practices for the publication of scholarly and research books. We are committed to the highest editorial standards in ethics and deontology, under the premise of "Open Science in Latin America and the Caribbean".



**OASPA**

**Editorial Mar Caribe**

**Chemical and food processes applied to goat's milk: Non-  
Conventional Derivatives**

**Colonia del Sacramento, Uruguay**

## About the authors and the publication

**Eduardo Zorrilla Tarazona**

[ezorrilla@unab.edu.pe](mailto:ezorrilla@unab.edu.pe)

<https://orcid.org/0000-0001-9038-7146>

*Universidad Nacional de Barranca, Peru*

**Juan Luis Pérez Marín**

[jperez@unia.edu.pe](mailto:jperez@unia.edu.pe)

<https://orcid.org/0000-0002-3671-1782>

*Universidad Nacional Intercultural de la  
Amazonía, Peru*

**Ingrid Yossy Robles Castañeda**

[iroblesc@unia.edu.pe](mailto:iroblesc@unia.edu.pe)

<https://orcid.org/0000-0001-6018-3644>

*Universidad Nacional Intercultural de la  
Amazonía, Peru*

**Ader Martín Guerra Rios**

[ader\\_guerra@unu.edu.pe](mailto:ader_guerra@unu.edu.pe)

<https://orcid.org/0000-0003-2458-9334>

*Universidad Nacional de Ucayali, Peru*

**Raúl Tello Suarez**

[rtellos@unia.edu.pe](mailto:rtellos@unia.edu.pe)

<https://orcid.org/0009-0002-7670-0321>

*Universidad Nacional Intercultural de la  
Amazonía, Peru*

**José Antonio Luna García**

[jlunag@unia.edu.pe](mailto:jlunag@unia.edu.pe)

<https://orcid.org/0000-0001-6309-5957>

*Universidad Nacional Intercultural de la  
Amazonía, Peru*

**Carlos Ruíz Padilla**

[carlos\\_ruiz@unu.edu.pe](mailto:carlos_ruiz@unu.edu.pe)

<https://orcid.org/0000-0001-9452-5438>

*Universidad Nacional de Ucayali, Peru*

### Book Research Result:

Original and unpublished publication, whose content is the result of a research process carried out before its publication, has been double-blind external peer review, the book has been selected for its scientific quality and because it contributes significantly to the area of knowledge and illustrates a completely developed and completed research. In addition, the publication has gone through an editorial process that guarantees its bibliographic standardization and usability.

## Index

Introduction .....	8
Chapter I .....	11
Chapter 1: The Unique Composition of Goat Milk: Innovations in Dairy Processing Methods .....	11
1.1 Goat Milk vs. Cow and Other Animal Milks: A Comprehensive Nutritional Comparison.....	11
1.1.1 Comparative Analysis with Other Animal Milks.....	13
1.1.2 Health Benefits and Considerations of Goat Milk .....	14
1.2 Unlocking the Biochemical Potential of Goat Milk: A Deep Dive into Its Unique Components and Innovative Applications .....	17
1.2.1 Fat Content and Types .....	18
1.2.2 Protein Profile and Quality .....	18
1.2.3 Vitamins and Minerals .....	19
1.2.4 Unique Biochemical Compounds in Goat Milk.....	19
1.3 Applications of Goat Milk in Unconventional Derivatives.....	21
1.4 From Tradition to Innovation: The Evolution of Dairy Processing Methods Through History .....	23
1.4.1 Traditional Dairy Processing Methods .....	24
1.4.2 Industrial Advancements in Dairy Processing .....	25
1.5 Modern Dairy Processing Innovations.....	27
Chapter II.....	31
Biological Processes in Dairy Innovation and Cutting-edge Techniques .....	31
2.1 Explain the recent advancements in technology and methodology that enable the production of non-conventional derivatives .....	31
2.1.1 Innovations in Data Analytics .....	34
2.1.2 Impact of Artificial Intelligence on Derivative Creation .....	36
2.2 Introduction to Biological Processes: Define key biological processes such as fermentation, enzymatic reactions, and microbial biotechnology .	38
2.2.1 Applications of Fermentation in Food and Beverage Industries ....	40

2.2.2	Role of Enzymes in Biological Reactions .....	41
2.2.3	Applications for Enzymes in Biotechnology and Medicine .....	42
2.2.4	Microbial Biotechnology .....	43
2.2.5	Impact of Microbial Biotechnology on Agriculture and Health .....	44
2.3	Exploring Unconventional Dairy Derivatives: The Health Benefits of Kefir, Probiotic Cheese, and Lactose-Free Products .....	46
2.3.1	Probiotic Cheese: Enhancing Traditional Cheese with Probiotics..	48
2.3.2	Lactose-Free Products: Catering to Lactose Intolerance and Beyond .....	50
2.3.3	Comparing Lactose-Free Products to Traditional Dairy .....	51
Chapter III	.....	54
Non-conventional derivatives of goat's milk	.....	54
3.1	Whey .....	54
3.1.1	Types of whey .....	57
3.2	Vitamins, minerals and nutritional properties .....	63
3.2.1	Cottage cheese (Whey cheese).....	66
3.2.2	Fermented milks.....	69
Chapter IV	.....	75
Technology for the production of fermented milks	.....	75
4.1	Types of fermented milks.....	76
4.1.1	Fermented milks with thermophilic lactic acid bacteria .....	76
4.1.2	Fermented milks with mesophilic lactic acid bacteria .....	77
4.1.3	Fermented milks with lactic acid bacteria and yeasts.....	77
4.1.4	Fermented milks with lactic acid bacteria and moulds .....	77
4.2	Condensed milks .....	80
4.2.1	Technological process for the production of condensed milk .....	80
4.2.2	Butter.....	81
4.2.3	Stages of production of goat's milk butter .....	82

<b>4.3 Navigating the Regulatory Landscape and Ensuring Quality Assurance in Goat Milk Development</b> .....	83
<b>4.3.1 International Standards and Compliance</b> .....	85
<b>4.3.2 Quality Assurance Measures in Goat Milk Processing</b> .....	86
<b>4.4 Future Directions in Goat Milk Development</b> .....	88
<b>Conclusion</b> .....	92
<b>Bibliography</b> .....	95

## Introduction

Goat milk has been consumed in many forms throughout history in many different cultures, prized for its nutritional profile and digestibility, and used directly and indirectly through products including cheeses, yogurts, and butter. Its unique matrix of protein structures and smaller fat globules make it a viable option for those who are intolerant to cow's milk. Goat milk is a dietary staple in many parts of the world and is praised not only for its versatility but also its potential health benefits.

Understanding these nutritional profiles is essential for consumers to make informed dietary choices. Factors such as protein quality, fat composition, vitamin and mineral content, and digestibility play significant roles in how each type of milk supports health. At the same time, the increasing awareness of lactose intolerance and milk allergies has prompted many to explore alternatives to conventional cow milk, leading to a rise in the popularity of goat, sheep, and other animal milks.

When comparing buffalo milk to goat milk, the latter again shines in terms of digestibility. Goat milk's lower fat content and different protein composition make it more suitable for individuals who may find buffalo milk too rich or difficult to digest. Yet more, goat milk's naturally occurring prebiotics can support gut health, which is an added benefit for many consumers. While buffalo milk may offer more fat and protein, goat milk provides a lighter, potentially more digestible alternative.

Instantaneous, the unique biochemical compounds found in goat milk, including short-chain fatty acids, bioactive peptides, and antioxidants, contribute significantly to its suitability for a range of unconventional derivatives. These compounds not only enhance its nutritional profile but also open doors for innovative applications across various industries, from food production to health and wellness sectors.



In this book the authors provide a comprehensive review of the evolution of processing methods in the dairy industry. By examining traditional techniques, industrial advancements, and modern innovations, we aim to highlight the historical context that has shaped dairy processing as we know it today. This exploration will offer insights into the ongoing developments and future trends that continue to influence this vital sector of the food industry.

In this sense, the general principle of processing methods in goat's milk are the same as those used in cow's milk, which consist of reducing the pH and activity of water to prolong its shelf life. The acid gel of goat's milk is characterized by firmness and lower viscosity compared to cow's and sheep's milk. The foundation of quality assurance in goat milk processing begins with stringent hygiene and sanitation practices. Maintaining a clean environment is crucial for preventing the introduction of pathogens and contaminants into the milk supply. This involves regular cleaning and sanitization of equipment, milking facilities, and storage areas.

Throughout the book, four chapters are explored; chapter I discerns the biological, chemical and nutritional composition of goat's milk and industrialized processes for processing its derivatives. In chapter II, the impact of microbial biotechnology on agriculture and health and the role of enzymes in biological reactions; in chapter III, non-conventional goat milk derivatives from artisanal to mass production are discussed and various chemical-industrial techniques and methods for obtaining lactose proteins, such as ultrafiltration, are envisioned. The use of acids as catalysts in protein precipitation and the use of heat treatments have also been feasible, the latter being the oldest process used for recovery. Chapter IV outlines a detailed treatment of lactic ferments and compilations of international standards, e.g. organizations such as the Codex Alimentarius Commission, established by the World Health Organization (WHO) and the Food and Agriculture Organization (FAO).

Therefore the research objective: To examine the chemical, biological and industrial processes for the synthesis of pasteurized goat milk and

conventional or unconventional derivatives, with emphasis on FAO and WHO regulations and recommendations. Therefore, it is important that the cheese industry has a portfolio of options to use whey as a food base, preferably for human consumption, in order not to pollute the environment and to recover, by far, the monetary value of whey, with the manufacture of whey powders, concentrated sweetener syrups for the food industry. The authors recommend the scientific community to complement the research presented in this book with local norms and regulations according to the type of derivative to be processed from goat milk and emphasize the importance of the agro-industrial processes presented in the book.

# Chapter I

## Chapter 1: The Unique Composition of Goat Milk: Innovations in Dairy Processing Methods

### 1.1 Goat Milk vs. Cow and Other Animal Milks: A Comprehensive Nutritional Comparison

Animal milk has long been a fundamental component of human diets across cultures and continents, prized not only for its taste likewise for its rich array of nutrients. The consumption of milk from various animals—such as cows, goats, sheep, buffalo, and camels—offers a diverse spectrum of nutritional benefits, each with unique profiles to cater to different dietary needs and preferences.

Understanding these nutritional profiles is essential for consumers to make informed dietary choices. Factors such as protein quality, fat composition, vitamin and mineral content, and digestibility play significant roles in how each type of milk supports health. At the same time, the increasing awareness of lactose intolerance and milk allergies has prompted many to explore alternatives to conventional cow milk, leading to a rise in the popularity of goat, sheep, and other animal milks (Walzem et al., 2002).

When comparing the nutritional profiles of goat milk and cow milk, several key factors must be considered, including protein content and quality, fat composition and digestibility, as well as the vitamins and minerals present in each type of milk. This analysis highlights the distinct advantages and characteristics of goat milk in relation to cow milk.

Both goat milk and cow milk are excellent sources of protein, essential for growth, repair, and overall health. Still, the protein structure in goat milk differs significantly from that of cow milk. Goat milk contains a higher proportion of short and medium-chain fatty acids, enhancing protein digestibility (Bendtsen et al., 2013). Therefore, the casein proteins in goat

milk form a softer curd compared to the firmer curd formed by cow milk proteins. This characteristic may contribute to easier digestion, particularly for individuals with sensitive stomachs.

In terms of overall protein content, cow milk typically has a slight edge, providing approximately 8 grams of protein per cup, while goat milk intentions around 7 grams. Nonetheless, the bioavailability of protein in goat milk may be more advantageous for certain individuals, especially those with gastrointestinal sensitivities.

Fat content is another critical area of comparison between goat milk and cow milk. Goat milk has a higher fat content, averaging around 4.5% to 5% fat compared to 3.5% to 4% in cow milk. Although, the primary type of fat in goat milk is medium-chain triglycerides (MCTs), which are more easily absorbed and metabolized by the body. MCTs have been associated with various health benefits, including improved energy levels and weight management.

Yet more, the fat globules in goat milk are smaller and more easily digestible than those found in cow milk. This can be particularly beneficial for individuals who experience difficulty digesting dairy products, as smaller fat globules may lead to less gastrointestinal discomfort and enhanced nutrient absorption.

Goat milk is known for its rich nutrient profile, containing a variety of essential vitamins and minerals. It is particularly high in calcium, phosphorus, and potassium, which are vital for bone health and overall bodily functions. Goat milk also contains higher levels of certain vitamins, such as vitamin A and vitamin B6, compared to cow milk.

On the other hand, cow milk is often fortified with vitamin D in many countries, crucial for calcium absorption and bone health. While goat milk does not naturally contain as much vitamin D, it recommends a wealth of other nutrients that contribute to a balanced diet. In sudden, while both goat milk and cow milk provide essential nutrients, goat milk distinguishes itself through its unique protein structure, digestibility, and

an array of vitamins and minerals. Understanding these differences can help consumers make informed choices based on their dietary needs and preferences.

### **1.1.1 Comparative Analysis with Other Animal Milks**

In the realm of dairy products, goat milk often garnishes attention for its unique nutritional profile and potential health benefits. Anyway, it is essential to compare goat milk not only with cow milk not only with other types of animal milks, including sheep, buffalo, and camel milk. This analysis will shed light on the distinct advantages and disadvantages of these various milks, helping consumers make informed dietary choices.

Sheep milk is another popular alternative to cow milk, particularly in Mediterranean diets. When comparing sheep milk to goat milk, the fat content is a notable difference; sheep milk contains a higher fat percentage, contributing to its creamier texture and richer flavor. Nutritionally, sheep milk is also higher in vitamins A, B12, and D, as well as calcium and zinc, making it a potent source of essential nutrients (Moatsou and Sakkas, 2019).

Although, goat milk has its own advantages. It tends to be easier to digest due to smaller fat globules and a different protein structure, which can be beneficial for individuals sensitive to cow milk. Also, goat milk contains higher levels of certain beneficial fatty acids, such as capric and caprylic acids, which may have antimicrobial properties. While both milks are nutritious, the choice between sheep and goat milk often comes down to personal taste and dietary requirements.

Buffalo milk is another alternative that stands out for its rich nutritional profile. It contains about 50% more fat than cow milk, contributing to its creamy consistency and making it ideal for producing rich dairy products like mozzarella cheese. Buffalo milk is also higher in protein and calcium, making it a nutrient-dense option for those looking to boost their intake of these essential nutrients (Bendtsen et al., 2013).

When comparing buffalo milk to goat milk, the latter again shines in terms of digestibility. Goat milk's lower fat content and different protein composition make it more suitable for individuals who may find buffalo milk too rich or difficult to digest. Yet more, goat milk's naturally occurring prebiotics can support gut health, which is an added benefit for many consumers. While buffalo milk may offer more fat and protein, goat milk provides a lighter, potentially more digestible alternative.

Camel milk has gained attention in currently for its unique health benefits and nutritional profile. It is lower in fat than both cow and goat milk while being rich in vitamins and minerals, particularly vitamin C and iron (Almasri et al., 2024). As a choice, the most significant advantage of camel milk is its potential therapeutic properties, particularly for individuals with diabetes. Research suggests that camel milk may help regulate blood sugar levels, making it an attractive option for those managing this condition.

In comparison to goat milk, camel milk is also noted for its hypoallergenic properties, containing a different type of casein protein that may be less likely to trigger allergic reactions. However, it is important to note that camel milk can be more expensive and less widely available than goat or cow milk, which may limit its accessibility for some consumers.

In hasty, while goat milk holds its ground as a nutritious choice, it is essential to consider the benefits and drawbacks of other animal milks, such as sheep, buffalo, and camel milk. Each type of milk bids a unique set of nutritional characteristics and health benefits, allowing individuals to select the option that best aligns with their dietary preferences and health needs.

### **1.1.2 Health Benefits and Considerations of Goat Milk**

Goat milk has garnered attention not only for its unique flavor and versatility in culinary applications similarly to its potential health benefits. Understanding these benefits, along with the considerations surrounding its consumption, can help consumers make informed dietary choices. In

particular, the most notable advantages of goat milk are its potential suitability for individuals who are allergic to cow milk or who suffer from lactose intolerance. Goat milk contains a different protein structure than cow milk, particularly in regard to casein, which may result in fewer allergic reactions. Some studies suggest that the proteins in goat milk could be less allergenic, making it a viable alternative for those sensitive to cow milk proteins.

Goat milk has a lower lactose content compared to cow milk, although it is not lactose-free. Many people who are lactose intolerant find they can tolerate goat milk better than cow milk due to the smaller fat globules in goat milk, which may facilitate easier digestion. Still, individuals with severe lactose intolerance should approach goat milk cautiously and consult healthcare professionals before making dietary changes milks (Walzem, 2004).

The nutritional profile of goat milk makes it particularly appealing for certain populations, including children and the elderly. For children, especially those with allergies to cow milk, goat milk can provide a rich source of essential nutrients such as calcium, phosphorus, and vitamins A and D, which are crucial for growth and development.

For the elderly, the digestibility of goat milk is a significant benefit. As individuals age, their digestive enzymes may become less efficient, making it harder to break down certain foods. The smaller fat molecules and different protein structure in goat milk can ease digestion, potentially aiding in nutrient absorption. In turn, its high calcium content supports bone health, a critical concern for older adults.

Beyond its health benefits, goat milk is celebrated for its culinary versatility. It can be used in a variety of dishes, ranging from cheeses like chèvre and feta to yogurt and ice cream. The distinct flavor of goat milk can enhance recipes, providing a creamy texture and unique taste that appeals to many palates (Donkor et al., 2007). Goat milk products have gained popularity across various cuisines worldwide, particularly in Mediterranean, Middle Eastern, and African cultures. The growing trend

toward sustainable and alternative sources of dairy has besides fueled the demand for goat milk, as many consumers seek to incorporate fewer common milks into their diets.

Immediate, while goat milk presents several health benefits and culinary possibilities, it is essential to consider individual dietary needs and potential allergies. As always, consulting healthcare professionals when making significant dietary changes is advisable, especially for those with existing health conditions.

The comparative analysis of goat milk versus cow milk and other animal milks reveals distinct nutritional profiles that cater to varying dietary needs and preferences. Goat milk stands out with its higher protein content and superior digestibility, making it an excellent alternative for those who may struggle with cow milk. Its unique fat composition, rich in medium-chain fatty acids, aids in easier absorption and may contribute to various health benefits.

When examining the vitamin and mineral content, goat milk advances a robust profile of essential nutrients, including higher levels of calcium, phosphorus, and certain B vitamins compared to cow milk. This positions goat milk as a particularly beneficial option for individuals looking to enhance their nutrient intake. Including, comparing goat milk with other animal milks, such as sheep, buffalo, and camel milk, underscores goat milk's versatility and uniqueness (Almasri et al., 2024). Each type of milk has its own set of advantages; for instance, sheep milk is richer in fat and protein, while camel milk boasts unique immunological properties.

The health benefits of goat milk are especially relevant for individuals with lactose intolerance or milk allergies, as it tends to provoke fewer allergic reactions and is often better tolerated (Moatsou and Sakkas, 2019). Now then, its nutritional profile makes it an appealing choice for children and the elderly, who may require easily digestible sources of nutrition.



In the culinary realm, goat milk's distinct flavor and versatility have contributed to its growing popularity in various cuisines worldwide. From cheeses to beverages, their applications are broadening, reflecting a shift in consumer preferences toward alternative dairy options.

Overall, goat milk emerges as a commendable source of nutrition, offering unique advantages over cow and other animal milks. As dietary choices continue to evolve, understanding the nutritional insights of different types of milk will empower consumers to make informed decisions that best suit their health and lifestyle needs.

## **1.2 Unlocking the Biochemical Potential of Goat Milk: A Deep Dive into Its Unique Components and Innovative Applications**

Goat milk has garnered significant attention in recent years, not only for its nutritional benefits but also for its unique biochemical properties that differentiate it from cow's milk and other dairy products. As consumers increasingly seek alternatives to traditional dairy, understanding the biochemical components of goat milk becomes essential in exploring its potential for various unconventional derivatives.

Biochemical properties refer to the specific chemical compounds and structures present in a substance that contribute to its functional characteristics and health benefits. In the case of goat milk, these properties are influenced by factors such as the breed of the goat, its diet, and the conditions under which the milk is produced. This chapter delves into the distinct biochemical components of goat milk, highlighting how these attributes enhance its nutritional profile and make it a suitable candidate for innovative applications in food, cosmetics, and health products.

Goat milk is rich in essential nutrients, including a unique combination of fats, proteins, vitamins, and minerals. Anyway, its true potential lies in the presence of specific bioactive compounds that offer numerous health benefits. From short-chain fatty acids that promote digestive health to bioactive peptides that may possess antimicrobial

properties, goat milk is a treasure trove of biochemical constituents that can be harnessed for a variety of uses.

Goat milk, often regarded as a nutritious alternative to cow's milk, boasts a unique nutritional profile that is increasingly drawing attention from health-conscious consumers and food innovators alike. Its composition not only makes it palatable for many who are lactose intolerant. In the same way provides a wealth of nutrients essential for overall health. Understanding the nutritional composition of goat milk is crucial for exploring its potential in unconventional derivatives.

### **1.2.1 Fat Content and Types**

One of the defining characteristics of goat milk is its fat content, which typically ranges from 3.5% to 4.5%. This fat is composed of a higher proportion of medium-chain fatty acids (MCFAs) compared to cow's milk, making it easier to digest and metabolize. MCFAs, such as caprylic and capric acid, are known for their quick absorption and energy-boosting properties. Accordingly, the fat globules in goat milk are smaller and more uniformly dispersed, which contributes to its creamy texture and enhances the absorption of fat-soluble vitamins. The unique fat profile not only affects the sensory qualities of goat milk apart from it opens doors for innovative products, particularly in the realm of functional foods and beverages.

### **1.2.2 Protein Profile and Quality**

Goat milk is also distinguished by its protein composition, which includes a higher proportion of essential amino acids compared to cow's milk. The primary proteins in goat milk are casein and whey, with the former being more prevalent. Goat milk contains a specific type of casein known as A2 beta-casein, which has been associated with improved digestibility and reduced allergic reactions in some individuals. The whey proteins present in goat milk, such as lactoglobulin and lactalbumin, are rich in bioactive peptides that may offer health benefits, including antimicrobial and immunomodulatory effects. This high-quality protein

profile makes goat milk an attractive option for those seeking dietary sources of protein, especially in unconventional applications.

### **1.2.3 Vitamins and Minerals**

Goat milk is a rich source of vitamins and minerals that are essential for various bodily functions. It contains significant levels of calcium, magnesium, phosphorus, and potassium, which are crucial for maintaining bone health, cardiovascular function, and muscle contraction. In perspective, goat milk is an excellent source of vitamins such as riboflavin (B2), vitamin B12, and vitamin D (ALKaisy et al., 2023). These nutrients play vital roles in energy metabolism, red blood cell formation, and calcium absorption, respectively. The bioavailability of these vitamins and minerals in goat milk is also enhanced due to the presence of beneficial fats, which aid in their absorption. This nutrient-rich profile supports the notion that goat milk can serve as a foundation for creating a diverse range of unconventional derivatives, from fortified foods to dietary supplements.

In extraction, the nutritional composition of goat milk—characterized by its distinct fat content, high-quality protein profile, and abundant vitamins and minerals—positions it as a valuable resource for innovative applications. As we delve deeper into the unique biochemical compounds that set goat milk apart, it becomes evident that its nutritional attributes are not just beneficial for direct consumption similarly hold promise for the development of unconventional derivatives.

### **1.2.4 Unique Biochemical Compounds in Goat Milk**

Goat milk is not only valued for its nutritional benefits purely for its unique biochemical compounds that contribute to its versatility and appeal in various applications. Understanding these distinct components can provide insight into why goat milk is increasingly recognized for unconventional derivatives.

In precise, standout features of goat milk is its higher concentration of short-chain fatty acids (SCFAs) compared to cow milk. SCFAs, such as butyric acid and caproic acid, are known for their potential health benefits,

including antimicrobial properties and their role in gut health. These fatty acids are rapidly absorbed and metabolized by the body, making them a valuable source of energy. Even more, the presence of SCFAs can enhance the flavor profile of goat milk, making it an excellent candidate for specialty products in the culinary world. Their unique properties also pave the way for goat milk to be utilized in functional foods aimed at promoting digestive health.

Bioactive peptides derived from goat milk have garnered attention for their potential health-promoting properties. These peptides are produced during the digestion of milk proteins and have been shown to exhibit various biological activities, including antihypertensive, antimicrobial, and immunomodulatory effects. The unique protein composition of goat milk, particularly its casein and whey proteins, contributes to the bioavailability of these peptides (Bendtsen et al., 2013). As a result, goat milk can serve as a functional ingredient in health-focused products, such as dietary supplements and fortified foods, aimed at improving overall wellness.

Goat milk is also rich in antioxidants and immunoglobulins, which play crucial roles in supporting immune function and combating oxidative stress. Antioxidants, such as selenium, vitamin E, and various phenolic compounds, help protect cells from damage caused by free radicals, promoting overall health and longevity. Immunoglobulins, particularly IgG, are antibodies that can enhance the immune response and provide protection against infections. The presence of these compounds makes goat milk an attractive option not only for health-conscious consumers similarly for the development of functional foods and supplements aimed at enhancing immunity and reducing the risk of chronic diseases.

Instantaneous, the unique biochemical compounds found in goat milk, including short-chain fatty acids, bioactive peptides, and antioxidants, contribute significantly to its suitability for a range of unconventional derivatives. These compounds not only enhance its nutritional profile In the same way open doors for innovative applications

across various industries, from food production to health and wellness sectors.

### **1.3 Applications of Goat Milk in Unconventional Derivatives**

Goat milk's unique biochemical properties lend themselves to a variety of unconventional applications that extend beyond traditional dairy products. As consumers increasingly seek alternative sources of nutrition and wellness, goat milk has emerged as a versatile ingredient in several innovative domains.

Fermented products made from goat milk have gained popularity due to their enhanced digestibility and probiotic benefits. The fermentation process transforms the lactose in goat milk into lactic acid, which not only reduces lactose content among which enriches the milk with beneficial bacteria. Products such as goat milk yogurt and kefir are noted for their creamy texture and tangy flavor, appealing to a broad audience milks (Walzem, 2004). To the maximum, the presence of short-chain fatty acids in goat milk promotes gut health, making these fermented products particularly attractive for individuals with lactose intolerance or sensitive digestive systems. The probiotic strains found in these products also contribute to improved immune function, farther solidifying goat milk's role in health-conscious diets.

The skincare industry has increasingly turned its attention to goat milk due to its rich composition of vitamins, minerals, and bioactive compounds. Goat milk is naturally high in fatty acids and vitamin A, both of which are known for their moisturizing and rejuvenating properties. These attributes make goat milk an ideal base for creams, lotions, and soaps that cater to sensitive skin types. Also, the presence of antioxidants and immunoglobulins in goat milk can help protect the skin from environmental stressors while promoting a healthy complexion. Products incorporating goat milk appeal to consumers looking for natural and effective skincare solutions, farther expanding the market for goat-derived cosmetics.

The culinary world has also embraced goat milk as a versatile ingredient in innovative food products. Its distinct flavor profile and nutritional advantages allow for the creation of gourmet cheeses, desserts, and beverages that stand out in a saturated market. Goat milk cheeses—such as chèvre and feta—are celebrated for their unique taste and creamy texture, making them a sought-after choice for gourmet cooking. In this regard, the use of goat milk in desserts, such as ice creams and custards, proposes a novel twist on classic recipes, providing a dairy alternative that is both delicious and nutritious. As food technology advances, the potential for developing new goat milk-based products continues to grow, appealing to health-conscious consumers and culinary enthusiasts alike.

In brief, the applications of goat milk in unconventional derivatives are vast and varied, driven by its rich biochemical composition and health benefits. From fermented products that promote gut health to skincare formulations that nourish the skin, and innovative food items that excite the palate, goat milk is carving out a significant niche in multiple industries. As awareness of these unique applications increases, the future of goat milk derivatives looks promising, paving the way for besides exploration and innovation.

As we look toward the future, the potential for goat milk derivatives is vast and promising. The unique biochemical properties of goat milk, including its distinctive nutritional composition and the presence of unique biochemical compounds, position it as a versatile ingredient across various industries. The increasing consumer demand for natural, healthy, and functional foods highlights the need for innovative applications of goat milk.

The exploration of fermented products opens up exciting avenues for enhancing the flavor profiles and health benefits of goat milk. As probiotic-rich foods continue to gain popularity, goat milk-based yogurts, cheeses, and kefir offer an attractive alternative to traditional dairy (Sánchez et al., 2009). Their unique taste, combined with the health benefits

of goat milk, can help cater to health-conscious consumers and those with lactose intolerance.

In the realm of cosmetics and skincare, the nourishing properties of goat milk are garnering attention. With its natural moisturizing qualities and rich array of vitamins and antioxidants, goat milk is becoming an increasingly sought-after ingredient in natural beauty products. The trend toward clean beauty and transparency in ingredient sourcing positions goat milk as an appealing option for formulators aiming to create effective and gentle skincare solutions.

In synopsis, the innovative food product landscape is ripe for exploration with goat milk. From gourmet cheese varieties to dairy-free alternatives crafted from goat milk, the culinary applications are limited only by creativity. As chefs and food innovators experiment with goat milk, we may see a rise in unique flavor combinations and product offerings that appeal to diverse palates.

In this respect, the future of goat milk derivatives is bright, supported by a foundation of rich biochemical properties that enhance its appeal across multiple sectors. As research continues to unveil the multifaceted benefits of goat milk, its role in both traditional and unconventional applications will expand, satisfying the evolving demands of consumers and paving the way for sustainable and innovative practices in the food and cosmetic industries.

#### **1.4 From Tradition to Innovation: The Evolution of Dairy Processing Methods Through History**

Dairy processing methods are essential for transforming raw milk into a variety of safe and consumable products, playing a crucial role in food safety and preservation. The significance of these methods extends beyond mere production; they ensure that dairy products remain nutritious and free from harmful pathogens, which is vital for public health. As dairy products are staples in many diets worldwide, the techniques employed in their processing have evolved over the centuries,

reflecting advancements in science, technology, and changing consumer preferences.

Historically, dairy processing has undergone a remarkable transformation, from rudimentary techniques used by ancient cultures to the sophisticated methods utilized in contemporary dairy production. This evolution has been influenced by various factors, including agricultural practices, technological innovations, and an increasing understanding of microbiology and food science. Each advancement has contributed to improving product quality, extending shelf life, and enhancing food safety.

The purpose of this chapter is to provide a comprehensive review of the evolution of processing methods in the dairy industry. By examining traditional techniques, industrial advancements, and modern innovations, we aim to highlight the historical context that has shaped dairy processing as we know it today. This exploration will offer insights into the ongoing developments and future trends that continue to influence this vital sector of the food industry.

### **1.4.1 Traditional Dairy Processing Methods**

The journey of dairy processing begins with the collection of milk, a practice that dates back thousands of years. In ancient times, milk was harvested directly from domesticated animals such as cows, goats, and sheep. Nomadic tribes relied on simple techniques to collect milk, often using hollowed-out containers made from natural materials. The initial handling of milk was rudimentary, focusing primarily on hygiene practices learned through experience rather than scientific understanding. Milk was typically strained to remove impurities, and in many cultures, it was consumed raw or used immediately for various products.

In specific, the earliest methods of dairy preservation involved fermentation. Cultures of beneficial bacteria were introduced to milk, leading to the development of products such as yogurt and kefir (Vilaplana, 2015). This process not only extended the shelf life of milk but also enhanced its nutritional profile and digestibility. The use of natural



fermentation was prevalent in many cultures, as it was a reliable means of transforming milk into safe and palatable forms. These fermented products became staples in various diets worldwide, showcasing the ingenuity of traditional dairy processors who understood the importance of microbial action in food preservation.

Cheese making stands out as one of the most prominent significant advancements in traditional dairy processing. The earliest evidence of cheese production dates back over 7,000 years, with archaeological findings in regions such as Mesopotamia and the Indus Valley. The process of cheese making involves curdling milk using rennet, which could be derived from the stomach lining of young ruminants.

This method allowed for the transformation of liquid milk into a solid form, making it easier to store, transport, and consume. Over centuries, the art of cheese making evolved, leading to a remarkable variety of cheeses, each reflecting the unique geographic and cultural contexts of its production. Cheese has become not only a vital source of nutrition as a significant cultural symbol, often linked to regional traditions and practices (Scott, 1991).

In concise, traditional dairy processing methods laid the groundwork for the industry as we know it today. Through a combination of practical knowledge and cultural practices, early dairy processors developed techniques that enabled them to preserve and enhance the nutritional value of milk. These methods formed the foundation upon which modern dairy processing would later build, blending tradition with innovation to meet the changing needs of society.

#### **1.4.2 Industrial Advancements in Dairy Processing**

The dairy industry has undergone significant transformations over the past century, particularly through industrial advancements that have revolutionized processing methods. These changes have not only enhanced efficiency and safety but have also responded to the growing demands of a global consumer market.

As a choice, all pivotal advancements in dairy processing were the introduction of pasteurization in the 19th century, named after the French scientist Louis Pasteur. Initially developed to prevent spoilage in wine and vinegar, pasteurization was soon adapted for milk. This method involves heating milk to a specific temperature for a set period to kill harmful bacteria without compromising product quality. The widespread adoption of pasteurization dramatically improved food safety and significantly reduced the incidence of milk-borne diseases such as tuberculosis and brucellosis. As a result, pasteurized milk became a staple in households, changing public perceptions of milk consumption and solidifying it as an essential dietary component.

As the demand for milk and dairy products surged, so did the need for more efficient production methods. The advent of mechanization in the early 20th century transformed the dairy industry. Milking machines replaced traditional hand-milking techniques, allowing for faster and more hygienic milk collection. This innovation not only increased the volume of milk that could be harvested solely improved the welfare of dairy cows by reducing stress and discomfort associated with manual milking.

In addition to milking, mechanization extended to various stages of dairy processing, including pasteurization, homogenization, and packaging. Automated systems and conveyor belts streamlined the entire production line, minimizing labor costs and maximizing output. These advancements led to the establishment of large-scale dairy farms and processing plants, which could produce milk and dairy products on a much larger scale than ever before.

With the rise of industrialization in the dairy sector came the need for standardized processing techniques to ensure consistency and quality across products. The establishment of industry standards and regulations helped define acceptable practices in dairy processing, including sanitation protocols, ingredient specifications, and nutritional labeling. Organizations such as the U.S. Food and Drug Administration (FDA) and

the European Food Safety Authority (EFSA) played crucial roles in developing these standards.

Standardization not only improved product safety and quality likewise facilitated trade, allowing dairy products to be exported across borders with confidence. Consumers began to expect uniformity in taste and quality, leading to the creation of branded products that could guarantee these attributes. This shift paved the way for the commercialization of a wide variety of dairy products, from fluid milk to cheese, yogurt, and ice cream (Scott, 1991).

In brief, the industrial advancements in dairy processing have dramatically reshaped the industry. The introduction of pasteurization, the mechanization of production, and the establishment of standardized processing techniques have collectively enhanced food safety, increased efficiency, and provided consumers with a diverse array of high-quality dairy products. As we move forward, these historical advancements will continue to serve as a foundation for further innovations in the dairy sector.

## **1.5 Modern Dairy Processing Innovations**

The dairy industry has undergone remarkable transformations over the past few decades, driven by technological advancements and a growing emphasis on sustainability. These modern innovations not only enhance the efficiency and safety of dairy processing but also address the evolving demands of consumers and environmental challenges.

Modern milk processing has been revolutionized by innovative technologies that improve the quality and safety of dairy products. Automated systems and advanced sensors are now employed to meticulously monitor and control processing conditions. Technologies such as ultra-high-temperature (UHT) processing allow milk to be heated to extreme temperatures for a brief period, effectively killing harmful bacteria while extending shelf life without the need for refrigeration. Immediately, high-pressure processing (HPP) has emerged as a non-

thermal method that preserves nutrients and flavors while ensuring food safety. These innovations not only enhance product quality among which streamline production processes, reduce waste and energy consumption.

The growing awareness of environmental issues has led the dairy industry to adopt more sustainable practices. Modern dairy processing facilities are increasingly implementing energy-efficient technologies, such as anaerobic digesters, which convert waste into biogas for energy production (Huang, 2024). Water conservation initiatives are also pivotal, with many dairies employing advanced filtration and recycling systems to minimize water usage.

In this respect, the shift toward plant-based dairy alternatives has spurred innovation in processing methods, requiring new techniques to produce high-quality products that meet consumer preferences while reducing the environmental footprint. The integration of sustainable practices not only helps mitigate the ecological impact of dairy processing likewise aligns with the values of a more environmentally conscious consumer base.

Looking ahead, the dairy industry is poised to embrace besides innovations that will shape its evolution. One significant trend is the increasing integration of artificial intelligence (AI) and machine learning in processing operations. These technologies can analyze vast amounts of data to optimize production efficiency, predict maintenance needs, and enhance quality control.

Now then, as consumers continue to seek healthier and more functional dairy products, research into fortification and probiotic enhancements will drive new processing methods. The focus on transparency and traceability in the supply chain is also expected to grow, with advancements in blockchain technology enabling consumers to access detailed information about the origin and processing of their dairy products.

Indeed, modern dairy processing innovations are characterized by technological advancements, sustainability practices, and a forward-looking approach that anticipates future consumer needs. As industry navigates these changes, the ability to adapt and innovate will be crucial for maintaining relevance and ensuring the continued success of dairy processing in a rapidly evolving marketplace.

The journey of dairy processing has been marked by significant advancements that have transformed how milk and its by-products are produced, preserved, and consumed. From the traditional methods of milk collection and fermentation to the introduction of pasteurization and mechanization, each phase of development has contributed to enhanced food safety and the longevity of dairy products. The historical significance of cheese making illustrates the ingenuity of early civilizations in utilizing available resources, while modern innovations have further refined these practices to meet the demands of a growing population (Scott, 1991). The evolution of dairy processing methods highlights a continuous pursuit of efficiency, quality, and safety, which is critical for both producers and consumers.

As we look to the future, the dairy industry remains in a state of flux, adapting to new complaints and opportunities. The integration of advanced technologies such as automation, artificial intelligence, and data analytics is revolutionizing production processes, enhancing quality control, and facilitating better supply chain management. Even more, the growing emphasis on sustainability is prompting industry to explore eco-friendly practices, such as waste reduction and energy-efficient processing methods. These ongoing changes not only reflect a response to consumer preferences but also align with global efforts to mitigate climate change and promote environmental stewardship.

The dairy industry stands at a crossroads where tradition meets modernity. The importance of innovation cannot be overstated; it is essential for the industry's survival and growth in an increasingly competitive market. As dairy processing methods continue to evolve,

stakeholders must embrace new technologies while preserving the cultural and historical significance of traditional practices. The future of dairy processing will be characterized by a balance between efficiency and sustainability, ensuring that industry can meet the demands of both today's consumers and future generations. In this dynamic landscape, the commitment to innovation will be paramount, driving improvements in quality, safety, and environmental impact, all of which are vital for the continued success of the dairy sector.

## Chapter II

### Biological Processes in Dairy Innovation and Cutting-edge Techniques

#### 2.1 Explain the recent advancements in technology and methodology that enable the production of non-conventional derivatives

In the ever-evolving landscape of financial markets, the term "*non-conventional derivatives*" has gained prominence as a descriptor for innovative financial instruments that diverge from traditional derivatives such as options and futures. These derivatives often feature unique underlying assets, structures, or payoff mechanisms that cater to the specific needs of investors and traders looking to hedge risk or speculate on market movements.

Non-conventional derivatives encompass a broad range of financial instruments that include exotic options, structured products, and hybrid derivatives, tailored to meet distinct investment objectives. Unlike standardized derivatives that are typically traded on exchanges, non-conventional derivatives are often customized and traded over-the-counter (OTC), allowing for increased flexibility in design and execution. This customization can involve various underlying assets, including commodities, cryptocurrencies, and even intangible assets like carbon credits, illustrating the diverse applications of these instruments within contemporary financial strategies.

The significance of non-conventional derivatives in financial markets cannot be overstated. They play a crucial role in risk management, enabling participants to hedge against specific risks that may not be adequately addressed by traditional derivatives. Including, these instruments can enhance market liquidity and facilitate price discovery by allowing for more tailored exposures to risk factors. As investors

increasingly seek to navigate complex market environments, the ability to create bespoke financial solutions has made non-conventional derivatives an essential component of modern portfolio management.

Recent advancements in technology and methodology have revolutionized the production and utilization of non-conventional derivatives. The integration of machine learning, big data analytics, and blockchain technology, among others, has improved pricing and risk assessment while streamlining the processes involved in creating and managing these instruments.

The financial landscape is continuously evolving, with sophisticated modeling techniques playing a crucial role in the creation and management of non-conventional derivatives. These advancements are pivotal not only for pricing and risk assessment not only for enhancing the overall efficiency and accuracy of financial operations. Below, we delve into some of the most significant advancements in financial modeling techniques that are shaping the industry (Kalogiannidis et al., 2024).

Currently, machine learning (ML) has emerged as a transformative technology in financial modeling, particularly in the realm of risk assessment. Traditional risk assessment methods often rely on historical data and predefined mathematical models, which can be limiting. However, machine learning algorithms can analyze vast datasets to identify patterns and correlations that might not be apparent through conventional analyses.

These algorithms adapt and improve over time as they process new information, allowing for more dynamic risk assessments. For instance, ML models can predict the likelihood of default or the potential impact of market shocks on derivative portfolios. By leveraging techniques such as neural networks and decision trees, financial institutions can enhance their predictive capabilities, leading to better-informed decision-making and improved risk management strategies.



Stochastic calculus has long been a cornerstone of financial modeling, particularly in the pricing of derivatives. Recent advancements in this mathematical framework have significantly enhanced its applicability to non-conventional derivatives. Stochastic calculus allows for the modeling of random processes, which is essential in a market characterized by volatility and uncertainty.

New methodologies, such as stochastic volatility models and jump diffusion processes, have been developed to capture the complexities of asset price movements. These models enable practitioners to better account for the inherent risks associated with non-conventional derivatives, leading to more accurate pricing and valuation (Webb, 2024). In theory, the integration of stochastic calculus with computational techniques has facilitated the development of robust algorithms that can process real-time data, further refining the pricing mechanisms for these financial instruments.

Monte Carlo simulations have gained prominence as a powerful tool for pricing complex derivatives, including non-conventional types. This computational technique relies on repeated random sampling to obtain numerical results, making it particularly useful in scenarios where analytical solutions are difficult or impossible to derive.

Recent advancements in Monte Carlo methods, such as variance reduction techniques and parallel computing, have significantly increased the efficiency and accuracy of these simulations. By utilizing high-performance computing resources, financial analysts can simulate a multitude of scenarios and assess the impact of various factors on derivative pricing. This level of analysis is crucial for understanding the potential risks and rewards associated with non-conventional derivatives, enabling traders and risk managers to make more informed decisions.

In synthesis, advancements in financial modeling techniques, particularly through machine learning, stochastic calculus, and Monte Carlo simulations, are revolutionizing the assessment and pricing of non-conventional derivatives. As these technologies continue to develop, they

will undoubtedly play a pivotal role in shaping the future of finance, allowing for more sophisticated and responsive financial instruments in an ever-changing market landscape.

### **2.1.1 Innovations in Data Analytics**

As the financial landscape evolves, the use of data analytics has become increasingly pivotal in the production and management of non-conventional derivatives. The integration of advanced analytical techniques allows market participants to gain deeper insights into market dynamics, enhance trading strategies, and improve risk management.

The rise of big data has transformed the way financial analysts approach market analysis. With the ability to process vast amounts of structured and unstructured data from diverse sources—such as social media, news articles, and economic indicators—market participants can uncover patterns and trends that were previously overlooked. Techniques such as data mining, predictive analytics, and machine learning algorithms enable analysts to identify correlations and forecast market movements with greater accuracy.

Even more, big data analytics facilitate the development of sophisticated trading algorithms that can respond in real-time to market changes. By harnessing predictive models, traders can optimize their strategies, thereby enhancing their ability to create and manage non-conventional derivatives that respond to unique market conditions.

Sentiment analysis has emerged as a powerful tool for understanding market psychology and its impact on asset prices. By analyzing textual data from social media platforms, financial news, and investor forums, traders can gauge public sentiment towards specific assets or market conditions. This qualitative data, when quantified, suggests insights that can be integrated into trading strategies, allowing for more informed decision-making.

For instance, if sentiment analysis indicates a growing negative perception of a particular asset class, traders can adjust their positions to

mitigate potential losses. Conversely, positive sentiment can signal investment opportunities. As sentiment analysis continues to evolve, its integration into trading algorithms will become more sophisticated, leading to more dynamic and responsive trading strategies for non-conventional derivatives.

The advent of blockchain technology has introduced revolutionary changes in the way financial transactions are recorded and verified. In the context of non-conventional derivatives, blockchain enhances data integrity and transparency, addressing some of the critical challenges associated with traditional derivative contracts.

By employing a decentralized ledger, blockchain ensures that all transactions are immutable and traceable, reducing the risks of fraud and counterparty default. This level of transparency fosters trust among market participants, encouraging more engagement in non-conventional derivatives. So, smart contracts—self-executing contracts with the terms of the agreement directly written into code—can automate the execution of derivative agreements, streamlining processes and reducing operational costs.

Concisely, the innovations in data analytics are instrumental in shaping the future of non-conventional derivatives, by leveraging big data techniques, sentiment analysis, and blockchain technology, market participants can enhance their analytical capabilities, optimize trading strategies, and ensure the integrity of their financial transactions (Magableh et al., 2024). As these technologies continue to evolve, their impact on the financial markets will be profound, driving further advancements in the production and management of non-conventional derivatives. The landscape of non-conventional derivatives is being transformed by a suite of emerging technologies that streamline production processes and enhance the efficiency and effectiveness of derivative creation.

### **2.1.2 Impact of Artificial Intelligence on Derivative Creation**

Artificial intelligence (AI) has revolutionized various sectors, and the financial markets are no exception. In the realm of non-conventional derivatives, AI algorithms can analyze vast datasets to identify patterns and correlations that human analysts may overlook. Machine learning models, in particular, enhance the ability to predict market movements and assess risks associated with different derivatives.

For instance, AI can automate the design of customized derivatives by leveraging historical data and current market conditions. This capability allows financial institutions to tailor products to specific client needs, creating unique offerings that drive market engagement. In this respect, AI-driven analytics can optimize pricing strategies, ensuring that derivatives are competitively priced while accounting for underlying risk factors.

Cloud computing has emerged as a critical infrastructure for the financial industry, particularly in the production of non-conventional derivatives. The ability to access vast computational power and storage on-demand allows financial institutions to scale their operations seamlessly. This scalability is crucial when developing complex derivatives that require extensive computational resources for modeling and analysis.

Cloud platforms facilitate collaboration among teams distributed across various locations, enabling real-time data sharing and communication. This is particularly advantageous for firms looking to innovate quickly and efficiently in response to market dynamics. Either, cloud computing enhances the security of financial data, which is essential given the sensitive nature of derivative contracts and the potential for cyber threats (Golightly et al., 2022).

The Internet of Things (IoT) is increasingly making its mark on financial markets, providing innovative solutions for monitoring and managing derivative activities. IoT technologies enable real-time data collection from various sources, including market feeds, economic

indicators, and even consumer behavior analytics. This influx of data can be instrumental in refining risk assessment models and enhancing decision-making processes.

For instance, IoT devices can track and report market conditions in real time, allowing traders to respond swiftly to changes that might affect the value of non-conventional derivatives. To the maximum, IoT integration can provide insights into the operational aspects of derivatives, such as trade execution and settlement processes. This transparency fosters greater trust among market participants and can lead to more efficient trading environments.

In hasty, the convergence of artificial intelligence, cloud computing, and IoT is reshaping the production processes of non-conventional derivatives. These technologies enhance efficiency and scalability while empowering financial institutions to innovate and respond to the evolving landscape of financial markets.

In this respect, the production of non-conventional derivatives has been profoundly influenced by a range of innovative technologies and methodologies. From machine learning algorithms that enhance risk assessment to sophisticated stochastic calculus applications and advanced Monte Carlo simulations, the advancements in financial modeling techniques have significantly improved our ability to price and manage these complex financial instruments. Including, innovations in data analytics, including big data methods for market analysis and sentiment analysis for trading strategies, have provided traders and financial institutions with deeper insights into market dynamics. The integration of blockchain technology has also revolutionized the integrity of derivative contracts, ensuring transparency and trust in transactions.

Emerging technologies are set to farther transform the landscape of non-conventional derivatives. Artificial intelligence is paving the way for more efficient derivative creation processes, while cloud computing extends the scalability necessary to handle the increasing complexity and volume of transactions. Therefore, the Internet of Things (IoT) is enhancing

the monitoring of derivative activities, allowing for real-time data collection and analysis.

Looking ahead, several potential challenges must be acknowledged. As technology continues to evolve, regulatory frameworks may struggle to keep pace with the rapid advancements, potentially leading to gaps in oversight and increased risks in the financial markets. In turn, reliance on technology could introduce vulnerabilities, such as cybersecurity threats, which need to be addressed proactively.

Future trends in non-conventional derivatives are likely to be characterized by greater customization and personalization in financial products, driven by the ability to analyze vast amounts of data and tailor offerings to specific investor needs. In theory, as markets become increasingly interconnected, the influence of global economic factors on derivative pricing and performance will become more pronounced, necessitating a holistic approach to risk management.

In the meantime, the intersection of technology and finance has ushered in a new era for non-conventional derivatives, presenting both opportunities and threat. As we advance, it will be crucial for market participants to stay abreast of these developments and adapt accordingly to navigate the evolving landscape effectively.

## **2.2 Introduction to Biological Processes: Define key biological processes such as fermentation, enzymatic reactions, and microbial biotechnology**

Biological processes are fundamental mechanisms that sustain life, enabling organisms to grow, reproduce, and adapt to their environments. These processes encompass a vast array of activities, from the cellular level to the intricate interactions within ecosystems. Understanding these processes is crucial, as they provide insights into how living systems function, interact, and can be manipulated for various applications in science and industry.

The significance of biological processes extends across multiple fields, including medicine, agriculture, environmental science, and biotechnology. In healthcare, for instance, a deep understanding of enzymatic reactions can lead to the development of more effective drugs and therapies. In agriculture, knowledge of fermentation processes enhances food production and preservation methods. Yet more, microbial biotechnology plays a pivotal role in sustainable practices, aiding in waste management and biofuel production. In this respect to exploring and harnessing these biological processes, their applications can lead to innovations that address all pressing ambitious facing humanity today.

Fermentation is a metabolic process that converts sugars into acids, gases, or alcohol in the absence of oxygen. This process is primarily carried out by microorganisms such as yeasts and bacteria, which utilize anaerobic pathways to extract energy from organic compounds (Vilaplana, 2015). Fermentation serves as an alternative to aerobic respiration, allowing organisms to generate ATP (adenosine triphosphate) in low-oxygen environments. The fundamental principle behind fermentation is the breakdown of glucose through a series of enzymatic reactions, leading to the production of byproducts such as ethanol, carbon dioxide, and organic acids.

There are several distinct types of fermentation, each characterized by the specific microorganisms involved and the end products generated. The two most notable forms of fermentation are:

- a. Alcoholic Fermentation:** This type is primarily conducted by yeast, particularly (*Saccharomyces cerevisiae*). During alcoholic fermentation, glucose is converted into ethanol and carbon dioxide. This process is widely utilized in the production of alcoholic beverages like beer and wine, as well as in baking, where the carbon dioxide produced causes dough to rise.
- b. Lactic Acid Fermentation:** This form of fermentation is performed by lactic acid bacteria such as (*Lactobacillus*). In lactic acid fermentation, glucose is metabolized as the main byproduct. This

process is essential in the production of yogurt, sauerkraut, and other fermented foods. It is also relevant in muscle metabolism, where lactic acid accumulates during intense exercise when oxygen levels are low.

Other types of fermentation include acetic acid fermentation, which is involved in vinegar production, and propionic acid fermentation, which plays a role in the ripening of certain cheeses. Each type of fermentation has unique characteristics and applications based on the microorganisms employed and the specific substrates utilized (Zhou and Sun, 2010).

### **2.2.1 Applications of Fermentation in Food and Beverage Industries**

The applications of fermentation in the food and beverage industries are vast and significant. Fermentation not only enhances the flavor, texture, and nutritional value of foods solely acts as a natural preservation method. The production of fermented foods has been practiced for centuries, with various cultures developing unique fermentation techniques.

In the beverage industry, fermentation is crucial for the production of alcoholic drinks. Beer production involves mashing grains to extract sugars, followed by fermentation using yeast, which converts these sugars into alcohol and carbon dioxide. Wine production similarly relies on the fermentation of grape sugars, with different yeast strains imparting distinctive flavors and aromas.

In the realm of food, fermentation contributes to the creation of probiotic-rich products. Foods such as yogurt, kimchi, and miso not only offer unique tastes apart from it promote gut health through the presence of beneficial microorganisms (Sánchez et al., 2009). Fermentation can also extend the shelf life of perishable items by inhibiting the growth of spoilage microorganisms.

In this vein, the rising consumer interest in health-conscious and naturally fermented products has fueled innovation in the food industry, leading to the development of new fermentation methods and products. As a result, fermentation remains a vital process that underpins both



traditional and modern culinary practices, reflecting its enduring significance in human culture and nutrition.

Enzymatic reactions are fundamental biological processes that catalyze chemical reactions within living organisms. Enzymes, which are typically proteins, function as biological catalysts that significantly accelerate the rate of these reactions without being consumed or permanently altered in the process (Bendtsen et al., 2013). Understanding how enzymes operate and the factors that influence their activity is crucial for various applications in biotechnology, medicine, and industrial processes.

### **2.2.2 Role of Enzymes in Biological Reactions**

Enzymes play a pivotal role in facilitating biochemical reactions by lowering the energy required for these reactions to occur. Each enzyme is highly specific to its substrate—the molecule upon which it acts—allowing for precise control over metabolic pathways (Robinson, 2015). For example, the enzyme amylase catalyzes the breakdown of starch into sugars, while lactase is responsible for the hydrolysis of lactose into glucose and galactose. This specificity is essential for maintaining the delicate balance of metabolic processes within cells.

In addition to catalyzing reactions, enzymes regulate metabolic pathways by acting as checkpoints. In response to varying cellular conditions, enzymes can be activated or inhibited, thereby influencing the overall metabolic activity of the organism. This regulatory function is vital for processes such as energy production, DNA replication, and the synthesis of biomolecules. Several factors influence the activity of enzymes, including temperature, pH, substrate concentration, and the presence of inhibitors or activators.

- a. Temperature:** Enzymes typically have an optimal temperature range within which they function most efficiently. Higher temperatures can increase reaction rates to a point, but excessive heat can denature

enzymes, leading to loss of function. Conversely, low temperatures can slow down enzyme activity.

- b. pH:** Each enzyme has an optimal pH at which it operates best. Deviations from this pH can result in decreased activity or denaturation. For instance, pepsin, an enzyme found in the stomach, operates optimally in acidic conditions, whereas trypsin in the small intestine functions best in a more alkaline environment.
- c. Substrate Concentration:** The rate of enzymatic reactions increases with substrate concentration up to a certain point. Once all active sites of the enzyme molecules are occupied, the reaction rate levels off, reaching a maximum velocity ( $V_{max}$ ). This phenomenon is described by the Michaelis-Menten kinetics.
- d. Inhibitors and Activators:** Inhibitors are molecules that decrease enzyme activity, either by blocking the active site or altering the enzyme's structure. Conversely, activators enhance enzyme function. Understanding these interactions is essential for developing drugs and therapies that target specific enzymatic pathways.

### 2.2.3 Applications for Enzymes in Biotechnology and Medicine

Enzymes have found extensive application in various fields, particularly in biotechnology and medicine. In biotechnology, enzymes are utilized in processes such as biofuel production, where cellulases break down plant biomass into fermentable sugars. In the food industry, enzymes like proteases and lipases are employed to improve the texture and flavor of products.

In medicine, enzymes are crucial for diagnostic tests and therapeutic applications. To illustrate, enzymes such as glucose oxidase are used in blood glucose monitoring devices for diabetes management. Enzyme replacement therapies have been developed for genetic disorders caused by enzyme deficiencies, such as Gaucher's disease and Fabry disease.

In this regard, enzymes are pivotal in the development of biopharmaceuticals, where recombinant DNA technology allows for the

production of human enzymes for therapeutic use. This has led to significant advancements in treating conditions ranging from clotting disorders to certain cancers.

As a consequence, enzymatic reactions are integral to life and have diverse applications that extend beyond natural biological processes. Understanding how enzymes function and the factors that influence their activity is essential for harnessing their potential in various scientific and industrial fields.

#### **2.2.4 Microbial Biotechnology**

Microbial biotechnology refers to the utilization of microorganisms—such as bacteria, fungi, and viruses—to develop products and processes that benefit humanity (Zhou and Sun, 2010). This interdisciplinary field combines microbiology, molecular biology, biochemistry, and engineering to harness the metabolic capabilities of microbes for various applications. The scope of microbial biotechnology is vast and includes areas such as pharmaceuticals, environmental management, agriculture, and food production.

By manipulating microbial systems, scientists can enhance natural processes, develop new technologies, and create sustainable solutions to pressing global complaint, including food security and disease management. The techniques employed in microbial biotechnology are diverse and continually evolved, driven by advancements in genetic engineering and molecular biology:

- a. Genetic Engineering:** This involves the modification of microbial genomes to enhance desirable traits or introduce new functions. Techniques such as CRISPR-Cas9 allow for precise edits to DNA, enabling the development of strains with improved metabolic pathways for increased product yield.
- b. Fermentation Technology:** Fermentation processes can be optimized using microbial cultures to produce a variety of bioproducts, including antibiotics, enzymes, and biofuels.

Controlling parameters such as temperature, pH, and nutrient availability is crucial for maximizing productivity.

- c. **Bioprocessing:** This technique focuses on the design and execution of processes involving living cells or components derived from them. Bioprocessing is essential in the production of biopharmaceuticals, where specific microbes are cultured under controlled conditions to produce therapeutic proteins or vaccines (Bendtsen et al., 2013).
- d. **Metagenomics:** By analyzing genetic material obtained directly from environmental samples, metagenomics allows researchers to explore the diversity and functionality of microbial communities. This technique has significant implications for understanding ecosystem dynamics and discovering novel bioactive compounds.
- e. **Bioinformatics:** The integration of computational tools in microbial biotechnology helps in analyzing large datasets generated from genomic and proteomic studies. Bioinformatics aids in identifying potential microbial strains for application in various biotechnological processes.

### 2.2.5 Impact of Microbial Biotechnology on Agriculture and Health

Microbial biotechnology has made substantial contributions to both agriculture and health, addressing all critical objections faced by these sectors.

- a. **Agricultural Applications:** Microbial biopesticides and biofertilizers are increasingly used to promote sustainable farming practices. Beneficial microbes can enhance soil fertility, suppress plant diseases, and improve crop resilience to environmental stresses. For instance, nitrogen-fixing bacteria play a crucial role in enriching soil nutrient content, reducing the need for synthetic fertilizers and minimizing environmental impact (Vilaplana, 2015).
- b. **Health and Medicine:** The pharmaceutical industry has benefited from microbial biotechnology, particularly in the production of antibiotics, vaccines, and probiotics. Microorganisms are used to

produce life-saving medications, such as penicillin and insulin. Then, probiotics derived from beneficial bacteria support gut health and contribute to overall well-being.

- c. **Environmental Remediation:** Microbial biotechnology also plays a vital role in bioremediation, where specific microbes are employed to degrade pollutants in soil and water. This process can effectively clean up oil spills, heavy metal contamination, and other hazardous waste, contributing to environmental sustainability.

In synopsis, microbial biotechnology is a dynamic field that harnesses the power of microorganisms to drive innovation across agriculture, health, and environmental management. By understanding and manipulating microbial processes, researchers are paving the way for sustainable solutions to some of the world's most pressing issues.

In abridgement, biological processes such as fermentation, enzymatic reactions, and microbial biotechnology play a pivotal role in numerous aspects of life and industry. We have explored the fundamental principles of fermentation, highlighting its various types—such as alcoholic and lactic acid fermentation—and its extensive applications in the food and beverage industries. The significance of enzymatic reactions has also been underscored, illustrating how enzymes are crucial catalysts in biological systems, with their activity influenced by various factors and their applications ranging from biotechnology innovations to medical advancements.

Moreover, the realm of microbial biotechnology showcases the profound impact that microorganisms have on agriculture and health, driven by sophisticated techniques that harness their capabilities for beneficial purposes (Zhou and Sun, 2010). This field continues to expand, offering promising solutions to global defiance, including food security and disease management.

Looking ahead, the study of biological processes is set to evolve besides, fueled by advancements in technology and our increasing understanding of biological systems. Future research may uncover new applications and

enhance existing processes, leading to breakthroughs that could transform industries and improve quality of life.

Hence, the significance of biological processes cannot be overstated; they are integral to both natural ecosystems and human endeavors. As we deepen our understanding of these processes, we open doors to innovative applications that could shape the future of medicine, agriculture, and environmental sustainability.

### **2.3 Exploring Unconventional Dairy Derivatives: The Health Benefits of Kefir, Probiotic Cheese, and Lactose-Free Products**

Nowadays, the landscape of dairy consumption has evolved significantly, driven by an increasing awareness of health, dietary restrictions, and the quest for enhanced nutritional benefits. Unconventional dairy derivatives—such as kefir, probiotic cheese, and lactose-free products—have emerged as innovative alternatives that cater to diverse dietary needs and preferences.

As more individuals turn to these unconventional options, it is crucial to understand their significance in modern nutrition. For many, lactose intolerance poses a barrier to enjoying traditional dairy, prompting the development of lactose-free alternatives that allow everyone to benefit from the nutritional value of dairy products without discomfort (Facioni et al., 2020). Now then, the rise of probiotic-rich foods reflects a growing appreciation for gut health and the role of probiotics in supporting overall well-being.

In this regard, integrating these unconventional dairy derivatives into daily diets aligns with contemporary health trends that emphasize functional foods that offer health benefits beyond basic nutrition. By exploring the various options available, from the tangy taste of kefir to the creamy richness of probiotic cheese, consumers can make informed choices that enhance their diets and promote a healthier lifestyle.

Kefir is a tangy, effervescent fermented dairy product that has gained popularity worldwide for its distinctive flavor and numerous

health benefits. Originating from the Caucasus region, kefir has been consumed for centuries and is recognized for its probiotic properties, which contribute to gut health and overall well-being.

Kefir is a cultured milk beverage produced by fermenting cow, goat, or sheep milk using kefir grains—small, gelatinous clusters of bacteria and yeast. These grains are rich in probiotics, which are live microorganisms that provide health benefits when consumed in adequate amounts (Zhou and Sun, 2010). The fermentation process typically takes 12 to 24 hours, during which the lactose in the milk is converted into lactic acid, giving kefir its characteristic tart flavor and creamy texture.

The origins of kefir date back to the Caucasus Mountains, where it was traditionally made by pouring milk into a leather pouch containing kefir grains. The unique environment and culture of the region contributed to the development of this probiotic-rich beverage. Today, kefir is enjoyed in various forms across the globe, from plain to flavored varieties, and is increasingly recognized for its health-promoting properties.

Kefir is not only a delightful addition to diet, not only a nutrient-dense food. It is an excellent source of protein, calcium, and vitamins such as B12 and riboflavin. Accordingly, kefir contains a wide variety of probiotics, which can enhance gut health by promoting a balanced microbiome.

The health benefits of kefir extend beyond its nutritional content. Research has shown that regular consumption of kefir may improve digestive health, boost the immune system, and even reduce inflammation. The probiotics found in kefir can aid in digestion, alleviate symptoms of lactose intolerance, and enhance nutrient absorption. At the same time, kefir is associated with potential benefits for mental health, as emerging studies suggest a link between gut health and mood regulation. Incorporating kefir into your diet is simple and versatile. Here are a few ideas to enjoy this fermented dairy delight:

- a. **Smoothies:** Blend kefir with fruits, vegetables, and a spoonful of nut butter for a nutritious and creamy smoothie that packs a probiotic punch.
- b. **Dressings and Dips:** Use kefir as a base for salad dressings or as a dip ingredient, mixing it with herbs, spices, and garlic for a refreshing twist.
- c. **Breakfast Bowls:** Pour kefir over granola or oatmeal, adding fresh fruits and nuts for a wholesome breakfast option.
- d. **Baking:** Substitute kefir for buttermilk or yogurt in baking recipes, such as pancakes, muffins, or cakes, to achieve a tender texture and enhance flavor.
- e. **Sipping:** Enjoy kefir straight from the bottle as a refreshing beverage or use it in place of milk in your favorite recipes.

This way, kefir is a versatile and nutritious fermented dairy product that poses a range of health benefits. Its unique flavor and probiotic content make it a valuable addition to modern diets, promoting better digestion and overall wellness. As we explore unconventional dairy derivatives, kefir stands out as a powerhouse of nutrition and a delightful culinary ingredient.

### **2.3.1 Probiotic Cheese: Enhancing Traditional Cheese with Probiotics**

As consumers become increasingly health-conscious, the food industry has responded with innovative products that merge traditional favorites with modern nutritional science. One such innovation is probiotic cheese, which incorporates beneficial live bacteria into the cheese-making process, enhancing both its flavor and health benefits. Probiotic cheese comes in various forms, offering options to suit different palates and culinary applications (Araujo et al., 2024):

- a. **Soft Cheeses:** Varieties such as cream cheese, ricotta, and feta often come with added probiotics. Their creamy textures and mild flavors make them versatile in both savory and sweet dishes.



- b. Hard Cheeses:** Cheeses like cheddar and gouda can also be produced with probiotics. These cheeses typically have a longer aging process, enhancing their complex flavors while still delivering the benefits of probiotics.
- c. Spreadable Cheeses:** Many brands offer probiotic-enhanced spreadable cheeses, ideal for dipping or spreading on crackers and bread. These varieties often combine flavors such as garlic, herbs, or spices.
- d. Plant-Based Probiotic Cheeses:** With the rise of veganism, plant-based dairy alternatives have emerged. Made from nuts, soy, or coconut, these cheeses can also be infused with probiotics, appealing to those seeking non-dairy options.

The incorporation of probiotics into cheese not only diversifies the flavor profile apart from it offers several health benefits:

- a. Gut Health:** Probiotic cheese contains live cultures that help balance gut microbiota, crucial for digestive health. This can lead to improved digestion and reduced gastrointestinal issues.
- b. Immune Support:** Regular consumption of probiotic-rich foods, including cheese, may enhance the immune system by promoting beneficial bacteria and inhibiting harmful pathogens.
- c. Nutritional Value:** Probiotic cheese retains the nutritional benefits of traditional cheese, including protein and calcium, while also adding the functional benefits of probiotics.
- d. Enhanced Flavor:** The presence of probiotics can contribute to unique flavor profiles, enriching the cheese experience. Many consumers find probiotic cheeses to have a more complex taste compared to their non-probiotic counterparts.

Probiotic cheese can be seamlessly integrated into various dishes, making it a delightful addition to any meal plan. Here are some creative ways to incorporate it:

- a. **Cheese Platters:** Include probiotic cheeses on charcuterie boards alongside fruits, nuts, and crackers for a delicious and nutritious appetizer.
- b. **Salads:** Crumble probiotic feta or add sliced probiotic cheddar to salads for a flavorful protein boost.
- c. **Sandwiches and Wraps:** Spread probiotic cream cheese on whole-grain bread or tortillas, pairing it with fresh veggies and lean proteins for a satisfying lunch.
- d. **Cooking and Baking:** Use grated probiotic cheese as a topping for casseroles, pizzas, or baked pasta dishes to enhance flavor while reaping health benefits.
- e. **Snacks:** Enjoy probiotic cheese on its own or with whole-grain crackers and fresh fruit for a quick, healthy snack.

Incorporating probiotic cheese into your diet not only elevates your culinary experience apart from it supports your health in multiple ways. As the demand for functional foods continues to rise, probiotic cheese represents a delicious and beneficial trend in the world of dairy.

### **2.3.2 Lactose-Free Products: Catering to Lactose Intolerance and Beyond**

Lactose intolerance is a common condition affecting a significant portion of the global population. It occurs when the body lacks sufficient levels of lactase, the enzyme required to digest lactose, the sugar found in milk and dairy products. In response to this widespread issue, the dairy industry has developed a range of lactose-free products, which not only cater to those with lactose intolerance among which appeal to individuals seeking to reduce their dairy intake for various reasons, including health and dietary preferences.

Lactose-free dairy products are made by treating regular milk with enzyme lactase, which breaks down lactose into simpler sugars—glucose and galactose—that can be easily absorbed by the body. This process results in milk that retains the same taste and nutritional profile as regular milk but is devoid of lactose. The range of lactose-free products has

expanded significantly and now includes milk, yogurt, cheese, and even ice cream. These options allow those who experience discomfort after consuming traditional dairy to enjoy familiar flavors and textures without the associated gastrointestinal distress (Sánchez et al., 2009).

Lactose-free products offer several health benefits beyond merely alleviating symptoms of lactose intolerance. For individuals who are lactose intolerant, these products enable the consumption of essential nutrients found in dairy, such as calcium, vitamin D, and protein, without adverse effects. In perspective, lactose-free dairy products often have higher digestibility than their traditional counterparts, making them suitable for individuals with sensitive stomachs or digestive issues.

In this vein, lactose-free products can be beneficial for those managing their overall sugar intake. Since lactose-free milk and yogurt have lower levels of naturally occurring sugars compared to regular dairy, they can be a better choice for individuals monitoring their sugar consumption for weight management or other health concerns.

### **2.3.3 Comparing Lactose-Free Products to Traditional Dairy**

When examining lactose-free products in comparison to traditional dairy, several factors come into play. Nutritionally, lactose-free options closely mirror their conventional counterparts, offering similar levels of protein, calcium, and other key nutrients. Even so, some lactose-free products may have a slightly sweeter taste due to the presence of broken-down sugars, enhancing the flavor without adding extra sugars or calories.

In terms of availability, lactose-free products have become increasingly mainstream, with many grocery stores stocking a variety of options. This accessibility has made it easier for consumers to make informed choices that suit their dietary needs. Even more, lactose-free products often cater to a broader audience, including those who may not have lactose intolerance but prefer to limit their dairy intake or are looking for alternatives that are easier on the digestive system.

Consequently, lactose-free products serve as a vital component in the modern dairy landscape, providing a delicious and nutritious alternative for those with lactose intolerance while appealing to a broader audience seeking healthier options. With their growing availability and diverse range, lactose-free products are not only transforming how individuals approach dairy consumption but are also contributing to overall dietary health and wellness.

Nowadays, there has been a notable shift in consumer preferences towards unconventional dairy derivatives, driven by a growing awareness of health and nutrition. Products like kefir, probiotic cheese, and lactose-free options are not just trendy alternatives; they represent a significant evolution in how we approach dairy consumption. As more individuals become concerned about gut health, lactose intolerance, and the benefits of probiotics, these innovative dairy products are finding their way into mainstream diets.

Kefir, with its rich probiotic content, offers a delicious and nutritious way to enhance gut health and boost the immune system. Its versatility allows it to be consumed in various forms, from smoothies to salad dressings, making it an easy addition to daily meals. Similarly, probiotic cheeses present an exciting blend of flavor and functionality, enriching our traditional cheese options with added health benefits. These cheeses not only satisfy culinary cravings on the contrary support digestive health, appealing to both cheese lovers and health-conscious consumers alike.

Lactose-free products have also emerged as a vital category, providing essential nutrients without the discomfort associated with lactose intolerance. With an increasing number of people identifying as lactose intolerant, these products help ensure that everyone can enjoy the nutritional benefits of dairy without adverse effects (Facioni et al., 2020). By bridging the gap between dietary restrictions and nutritional needs, lactose-free options are enhancing the inclusivity of modern diets.

So, to embrace a more health-focused approach to eating, unconventional dairy derivatives will become even more prominent. Their

unique health benefits, combined with culinary versatility, make them valuable additions to our diets. In this era of informed eating, understanding and incorporating these innovative dairy products can lead to improved health outcomes and a more enjoyable eating experience. Thus, the relevance of unconventional dairy derivatives is not just a passing trend; it is a fundamental shift towards a healthier, more inclusive approach to nutrition in contemporary society.

## Chapter III

### Non-conventional derivatives of goat's milk

#### 3.1 Whey

Whey or whey is the liquid fraction obtained during the coagulation of milk in the process of making cheese and casein, after the separation of the clot, its characteristics correspond to a greenish-yellow liquid, cloudy, with a fresh taste, weakly sweet. Whey is one of the most polluting materials in the food industry. Every 1000 liters of whey about 35 kg of biological oxygen demand (BOD) and about 68 kg of chemical oxygen demand (COD) are generated. This polluting force is equivalent to that of sewage produced in one day by 450 people (FAO, 1980).

Whey is defined as a liquid obtained after the precipitation and separation of casein from milk during cheese production and constitutes 85% - 90% of the volume of milk (González 2010), whose main components such as lactose, calcium, mineral salts and low molecular weight lacto-serum proteins soluble at their isoelectric point are retained in 55%. since they do not react with rennet (Caro et al., 2011).

Whey is a highly diluted product and its organoleptic and physicochemical characteristics can vary depending on the type of whey (sweet or acidic) or what is the same type of cheese processing (Aider et al. 2009; Fernández et al. 2009), the source of milk (cow, goat, buffalo, sheep, etc.), the animal's diet, the time of year and the lactation status (Valencia 2009). Previously, this by-product was described as an environmental pollutant, since when it was discharged into the environment without any type of treatment to counteract the high content of protein and sugary components, it had an impact on the quality of the waters, generating high rates of BOD (Biochemical Oxygen Demand) and COD (Chemical Oxygen Demand) (FAO 2009).

After various investigations, the mentality of whey has changed, which is not a whole substitute for cow's milk because it is a fraction of it

but contains nutrients and compounds with potential nutritional and health benefits, making it an important component for the manufacture of other products. There is very little statistical information on whey production, which may be basically due to the lack of interest in it and its use, including the lack of interest in it as a raw (Zhou and Sun, 2010).

Not using whey as food is a huge waste of nutrients; Whey contains just over 25% of milk proteins, about 8% of fat and about 95% of lactose. As shown above, at least 50% of the nutrients in milk remain in the whey. The same 1000 liters of whey referred to above contain more than 9 kg of protein of high biological value, 50 kg of lactose and 3 kg of milk fat. This is equivalent to the daily protein requirements of about 130 people and the daily energy requirements of more than 100 people. Therefore, it is important that the cheese industry has a portfolio of options to use whey as a food base, preferably for human consumption, in order not to pollute the environment and to recover, by far, the monetary value of whey, with the manufacture of whey powders, concentrated sweetener syrups for the food industry. (FAO, 1980).

Goat's milk whey (LSLC) is the liquid resulting from the production of curd or cheese, when casein and fat are separated during coagulation (about 85-90% of the volume of milk used). It is made up of approximately 93% water, 5% lactose, a little less than 1% protein, of which half are proteins of high nutritional value such as; albumins, globulins and protease-peptone, 0.7% minerals, with a higher content of sodium, potassium, magnesium, chloride and phosphate, also contains the water-soluble vitamins of milk, the most important is riboflavin, it also has small amounts of fat and lactic acid (LA) and its pH is between 5-6.

Currently the serum has little industrial applicability, the surpluses become a very high pollutant (BOD), from 40000 to 60000 ppm and a chemical oxygen demand (COD) of 50000 to 80000 ppm. More than 90% of these demands are due to the transformation of lactose. Hence, the search for alternatives for the use of this effluent becomes a national need. One of the alternatives is to obtain lactic acid (Plata et al., 2013).

Lactic acid is widely used in industry. Due to its characteristics, it is used in the food industry (in beverages and as a preservative), in pharmacy, medicine, textiles, in the leather industry and for the production of biodegradable plastics. One of the main difficulties in large-scale production of LA is the cost of raw materials, and it is of interest to find new low-cost means to improve the economics of the process.

The major organisms that produce LA belong to the families Streptococcaceae (genera *Streptococcus*, *Lactococcus*, *Leuconostoc*, *Pediococcus*, *Aerobacter* and *Gemella*) and Lactobacillaceae (genus *Lactobacillus*). The Lactobacillaceae family and especially those of the genus *Lactobacillus* (*L. bulgaricus*, *L. helveticus*, *L. delbruekii*, etc.), are facultative anaerobes (they do not have surface growth or it is scarce), they do not have catalase, they do not reduce nitrate; its metabolism is fermentative, about 50% of its final product is lactic acid, its optimal growth temperature is between 30-40°C, in an acceptable range between 26 and 46°C, at a pH of 4.5-7.2 and it is a non-pathogenic microorganism (Plata et al., 2013).

This by-product of the cheese-making process retains about 55% of the nutrients of the milk, among which are serum proteins with an appropriate balance in amino acids, high digestibility and excellent functional characteristics, which has induced the development of fractionation processes and concentration of its constituents. In some countries, the application of fractionation and concentration processes in the treatment of whey has made it possible to obtain concentrates with a protein content between 30 and 80%, which are being used for the enrichment of foods such as bread, soups and beverages, helping to alleviate protein deficiencies caused by excessive population growth. and due to the scarcity and increase in the price of conventional protein foods (Tirado et al., 2015).

Whey has significant protein, fat, lactose and calcium content. In relation to protein content, the serum proteins in milk are globular, among them, those present in greater quantities are  $\beta$ lactoglobulin ( $\beta$ -LG) and  $\alpha$ -



lactalbumin ( $\alpha$ -LA) and as minor constituents are lactoferrin, lactoperoxidase, immunoglobulins and glycomacropetides, among others. Although whey, as an abundant and readily available by-product of the cheese industry, was once considered a waste material, it is now considered a valuable source of protein and is widely used as a food ingredient.

Various techniques and methods have been used to obtain lactoseic proteins, such as ultrafiltration. It has also been feasible to use acids as catalysts in protein precipitation and the use of heat treatments, the latter being the oldest process used for recovery. The food industry, and dairy in particular, has always been interested in finding new methods for food preservation in order to improve the hygiene and safety of the final product, increase its shelf life and maintain a natural flavor in the food (Tirado et al., 2015).

### **3.1.1 Types of whey**

Depending on the type of coagulation of the casein used in cheese manufacturing, sweet whey or acidic whey is generated. Sweet whey is obtained from enzymatic coagulation, with a pH close to that of fresh milk, which, due to the stability of its composition, is the most widely used in the industry, unlike acid whey, which results from acid or lactic coagulation. For Jelen (2003), there are several types of whey depending mainly on the elimination of casein, the first called sweet, is based on coagulation by renin at pH 6.5. The second so-called acid results from the process of fermentation or addition of organic acids or mineral acids to coagulate casein as in the production of fresh cheeses. Coagulation occurs by natural acidification or the addition of organic acids.

On the other hand, González (2010) defines sweet whey as that obtained by enzymatic coagulation using a rennet of animal origin, such as calf renin or a microbial rennet of genetic technology. Acid whey is obtained by the natural acidification of milk or by the fermentation of milk, due to the bacterial flora existing in it and that obtained by the addition of acids. Coagulation is mainly caused by chemical and/or bacterial

acidification. On the other hand, whey, or whey, sweet is that whey in which the lactose content is higher and the acidity is lower than that of acidic whey.

In summary, Pintado et al. (2012) define sweet whey as the aqueous phase that is separated from curd in the process of making cheese or casein, greenish-yellow in color with a pH between 5.8-6.6 and Riera et al. (2004) acid whey as that produced in dairy industries when coagulation is carried out with an acid, decreasing the pH value to 5.1. Each of these definitions are similar in their clarifications from their obtaining to the pH range that are used in each of them, which allows this concept to be clearly explained.

#### **a. Whey technology**

Regardless of the subsequent treatment, the first step is the separation of the fat and casein fines, as they interfere with the subsequent steps. Fine casein particles are always present in the whey. These have an adverse effect on the fat separation process, which is why they must be separated in the first place. The whey to be stored, before treatment, must be cooled or pasteurized as soon as the fat contained in it is separated. When storage is going to be short (10-15 hours), cooling is sufficient to reduce bacterial activity. If the whey is to be stored for longer periods, it is necessary to proceed with pasteurization (Kreczmann et al., 2015).

#### **b. Obtaining WPC**

Whey protein concentrates are presented as a powder manufactured by drying the ultrafiltration concentrate. They are described in terms of their protein content (percentage of protein over dry matter), ranging from 35 to 80%. To make a product with 35% protein, liquid whey is concentrated about six times to a total solids content of approximately 9%. To obtain a concentrate with 80% protein, the liquid whey is concentrated about 20-30 times by ultrafiltration to a solids content of approximately 25%. This value is considered as the maximum for an economical operation, so it is necessary to diafilter the concentrate to remove lactose and ash and increase the concentration of proteins in

relation to the total dry matter. Diafiltration (DF) is a process in which water is added to the food as filtration is carried out in order to wash away the low molecular weight components that will pass through the membrane, basically lactose and mineral salts.

This permeate is used as a raw material to obtain lactose. Before the ultrafiltration stage, additional treatments, such as demineralization by nanofiltration (NF), can be performed. This is done through the use of specially designed, very small-pore reverse osmosis membranes, where small particles such as certain monovalent ions (sodium, potassium, chloride) and small organic molecules (such as urea and lactic acid) can escape through the membrane along with the aqueous permeate to obtain a protein concentrate (WPC) from whey. Most of the pure proteins, usually more than 99%, are retained in the concentrate, along with the fat content that was not removed in skimming (Kreczmann et al., 2015).

### **c. Obtaining WPI**

Whey protein isolates can be obtained mainly by two methods: membrane technology and ion exchange technology. The exchange of ions provides a higher concentration of protein per kilo of final product, but this is only part of the equation that is of interest. Ion exchange isolates, like many whey concentrates, are processed at high temperatures and therefore sacrifice the biological activity of the micro fractions that exist in whey. These ion exchange isolates are obtained through a technology called an "ion exchange column", which separates proteins from the rest of the components of the serum by discriminating based on their electrical charge.

During this process, most of the bioactive components, which give whey protein extra value to promote health and performance, are broken down or lose their biological capacity altogether. One of the sharpest losses occurs in the so-called glycomacropptides (GMP). Instead, ion exchange isolates manage to increase (Kreczmann et al., 2015). The concentration of beta-lactoglobulin is alarming, a subfraction that is not at all interesting since its indiscriminate ingestion is associated with a large number of

allergic reactions. Membrane technology uses series stages of ultrafiltration (UF) and microfiltration (MF) to obtain WPI; i.e. the MF permeate (degreasing) is sent to a second UF unit for further concentration.

This stage also includes diafiltration (DF). The microfiltration membranes used for this purpose are designed to retain suspended particles in the micron range, such as fats present in concentrated whey. On the other hand, the ultrafiltration membranes used are designed to retain whey constituents in the molecular range, such as proteins, separating them from lactose and other impurities present in whey (raw material for the production of lactose).

It should be noted that treating whey concentrate from ultrafiltration in a microfiltration plant can reduce the fat content of WPC 80-85% powder from 7.2% to less than 0.4%. This technique differs from ion exchange in that there is no chemical modification of the proteins and that the glycomacropeptide fraction is retained along with the others. If there is no pH adjustment and the process is carried out at intermediate temperatures, the final product is almost completely free of denatured proteins (Kreczmann et al., 2015).

#### **d. Composition of whey**

The main components of whey, both sweet and acidic, are water (93-94%), lactose (70-75% of total solids), serum proteins (8-11% of total solids) and minerals (10-15% of total solids). It also has fat and vitamins. The composition of whey depends on the type of cheese (enzymatic or acidic), the cheese-making techniques used (such as the coagulation method), the treatment undergone by the liquid whey (heat treatments, pre-concentration, recovery of casein fines), the physiological state of the animal, the type of breed and species, and also follows the trend of the chemical composition of the milk from which it comes (Jelen, 1992).

#### **e. Lactose**

Lactose is not only the main carbohydrate in milk, but also the main component of whey (apart from water), constituting approximately 4.4 -

4.9% of total whey (~ 75% of dry extract) (Jelen, 1992). It is a disaccharide formed by a glucose molecule and a galactose molecule, which is synthesized in the mammary gland. The concentration of lactose in the serum is fairly constant, but it depends on the proportion of the original lactose that has been degraded to lactic acid. Acid whey has a lower lactose content than sweet whey, and consequently, a high lactic acid content because during bacterial fermentation part of the lactose is transformed into lactic acid. It is reducing sugar, which in some circumstances, can react with the amino groups of proteins.

It is an adequate source of energy that plays an important role in the absorption of calcium, lactose is characterized by having limited solubility and low sweetness which limits, on many occasions, its use in food, so it is usually hydrolyzed before use. Yellowing of serum may occur after prolonged heat treatment or even by non-enzymatic browning reactions (Maillard reaction) between lactose and serum proteins (Jelen, 1992).

#### **f. Serum proteins**

Although the main component after water is lactose, serum proteins are the most valuable ingredient (Jelen, 2003). They can be defined as proteins that remain soluble in the liquid phase after precipitation of caseins at pH 4. They make up approximately 0.7% of the whey (~8-11% of the dry extract). In addition, whey also contains 0.2-0.3% non-protein nitrogen. The protein content of whey depends mostly on the type of clot and its treatment, and the presence of curd particles in it can increase it considerably (Scott, 1991). In milk, 80% of the proteins are caseins; however, these are added to the curd during cheese formation, while serum proteins are not retained in the curd, remaining dissolved in the aqueous portion (Walzem et al., 2002).

Serum proteins are even a more heterogeneous group than caseins, and have few features in common (Robin et al., 1993), exhibiting a number of unique properties that depend on their molecular weight, composition, and amino acid sequence.

- They are soluble under conditions that are not caseins (at pH 4.6-4.7).
- Most are globular proteins.
- They are easily denatured by heat, while caseins are more stable.
- Caseins have strong hydrophobic regions while serum proteins have a greater balance between hydrophilic and hydrophobic residues.
- They are not greatly affected by pH and salts.
- They are not phosphorylated and are insensitive to calcium.
- They have a more organized secondary and tertiary structure than caseins due to a more uniform distribution of amino acid types along the polypeptide chain and the presence of disulfide bridges (which implies that they have large amounts of cysteine unlike what happens in caseins). All serum proteins contain intermolecular disulfide bridges that stabilize their structure.

One of the most important characteristics of serum proteins is the high presence of amino acid residues with sulfhydryl groups that allow them to form intermolecular covalent bonds during processes at high temperatures (which causes their denaturation and subsequent aggregation). Whey proteins are soluble over a wide pH range and are denatured (individually and in solution) between 64 and 85 °C, and from this temperature they begin to aggregate and gel. Denaturation involves the loss of protein solubility. The main serum proteins are  $\beta$ -lactoglobulin ( $\beta$ -LG),  $\alpha$ -lactalbumin ( $\alpha$ -LA), immunoglobulins (Igs), serum albumin (BSA) and peptone proteoses. Along with these, other minority proteins such as glycomacropeptide, lactoferrin, and enzymes such as lactoperoxidase appear. The differences found in the protein profile may be a consequence of seasonal variations, the stage of lactation, the breed, the diet and the cheese processing techniques used. The different heat treatments applied, both to milk and whey, can also influence serum

proteins, since their heat stability is different, and follows this decreasing order:  $\alpha$ -LA >  $\beta$ -LG > BSA > Ig (Morr and Ha, 1993).

#### **g. Lipids**

Goat's milk contains 97-99% free lipids and 1-3% lipids bound to other substances (Park, 2006). Free lipids are made up of triglycerides or triacylglycerols, diglycerides or diacylglycerols and monoglycerides or monoacylglycerols, while bound lipids are made up of neutral lipids (mainly triglycerides) and polar lipids (phospholipids and glycolipids).

Practically all milk fat is found in fat globules, with the average diameter of fat globules in goat's milk ( $2.76\mu\text{m}$ ) being smaller than those in cow's milk ( $3.51\mu\text{m}$ ). The small size of the globules implies a better dispersion and a more homogeneous mixture of the fat in the milk, although it has a worse ability to skim (Park, 2006). In addition, the fat globules in goat's milk do not experience coalescence at refrigeration temperatures due to the absence of agglutinin, which is responsible for the aggregation of fat globules in cow's milk. Lipids present in serum are associated with the membrane of the fat globule and phospholipids are the main lipids of this membrane (Walstra et al., 2006).

However, triacylglycerols are the main lipids in CPS, followed by phospholipids, diacylglycerols, free fatty acids, cholesterol esters, cholesterol and monoacylglycerols, which may be a consequence of the presence of small fat globules in the serum. Serum lipid concentration is low (Walzem, 2004). It is usually 0.5-1%, although it depends on the type of milk, the type of cheese and the efficiency of the cheese manufacturing process. In most sour cheeses such as cottage or quark, the lipid content is almost negligible because the type of milk used is usually skimmed. (Jelen, 1992).

### **3.2 Vitamins, minerals and nutritional properties**

Minerals make up the third largest component of the total solid content of whey. Although the salt content is usually quite constant, its mineral composition varies depending on the type of whey obtained and

depends on other factors related to the cheese manufacturing methods used, especially the addition of sodium chlorides or calcium chlorides to milk. However, the main differences between the two types of whey are due to the different way in which caseins coagulate. Acidic sera have a higher calcium and phosphorus content due to the solubilization of colloidal calcium phosphate from casein micelles during acidification.

However, in sweet whey, calcium is not separated from the micelles, so most of the calcium remains retained in the cheese and is not lost in the whey. Acidic whey is also richer in magnesium and zinc, and citrate. The main macro minerals in goat's milk are calcium, phosphorus, potassium and chlorine, with a higher content than in cow's milk, and above all, in relation to human milk (Sanmartín, 2010; Ballard and Morrow, 2013).

Calcium appears to be important in terms of heat stability, while sodium and potassium are responsible for the salty taste, and along with lactose for whey osmolarity. During the processing of the serum, the concentration of minerals can be altered, as its content can be reduced by electro dialysis, nanofiltration or ion exchange. During ultrafiltration (UF), minerals bound to proteins (e.g., calcium, phosphorus, magnesium) are retained in the membrane and concentrated together with them, while soluble minerals (sodium, potassium, citrate) pass into the permeate (Sanmartín, 2010).

Riboflavin (B2) is the most abundant vitamin in whey. It is a water-soluble, heat-stable, and light-sensitive protein. It is responsible for the greenish color of the serum. It can be retained by reverse osmosis, but not through UF membranes, so UF permeates can also have this color. Goat's milk, and therefore whey, has higher amounts of vitamin A than cow's milk, because the goat converts all  $\beta$ -carotene into vitamin A, while it is deficient in folic acid and vitamin B12 compared to cow's milk. Both milks are deficient in pyridoxine (B6), vitamin C and D. The water-soluble vitamins that are present in milk remain in whey in varying proportions: 40-70% of vitamin B12, 55-75% of B6 and pantothenic acid, 70-80% of riboflavin and biotin, 80-90% of thiamine, nicotinic acid, folic acid and



ascorbic acid. The concentration of vitamin C is reduced during processing, so whey is not considered a good source of this protein. Of all the fat-soluble vitamins, the most abundant is A. During UF, fat-soluble vitamins are retained in the concentrate by separating with lipids, while only folic acid and vitamin B12 are retained from soluble vitamins (Sanmartín, 2010).

Since ancient times, the nutritional value of whey has been recognized. In 460 BC, Hippocrates recommended drinking considerable amounts of serum for long periods, because he had observed that it had therapeutic applications (Jelen, 2003). During the Middle Ages, many doctors prescribed the serum against various diseases. In the mid-seventeenth century, whey became the fashionable drink in English cities, opening numerous "whey houses", analogous to today's "coffee shops". During the nineteenth century, serum baths were very popular in spas (Tunick, 2008); in the 40s in the spas of Central Europe many diseases, such as anemia, tuberculosis, arthritis, uremia, etc., were treated with the ingestion of up to 1.5 kg of serum/daily (Holsinger et al., 1974).

Whey is a good source of bioavailable nutrients, providing high-quality protein, lactose, minerals, water-soluble vitamins (especially riboflavin), and bioactive components. From a nutritional point of view, proteins are the main focus of interest because they are an excellent nutritional source, as they are easily digestible and contain all the essential amino acids in the right proportions and similar to the proteins of the human body. Serum proteins have the highest bioavailability of all proteins, so our body can metabolize them more efficiently (Pordesimo and Onwulata, 2008).

Serum proteins, unlike caseins, pass rapidly from the stomach to the small intestine during digestion, where they are hydrolyzed more slowly while their amino acids are absorbed more quickly than amino acids from caseins. For this reason, serum proteins can be used as emulsifiers in food formulation to decrease intestinal transit time, thereby improving nutrient digestibility and absorption.

They have a high biological value, compared to that of other dietary proteins, thanks to their high content of essential amino acids (mainly  $\alpha$ -LA and  $\beta$ -LG) and especially lysine, tryptophan, methionine and cysteine). The high intake of some essential amino acids (lysine, threonine, methionine and isoleucine) allows whey to be used as a supplement to vegetable proteins (Walzem, 2004).

Whey has bioactivity, which is attributed to serum proteins and other nitrogenous compounds (bioactive peptides), which are those protein fragments obtained by proteolysis after the action of gastrointestinal enzymes. For peptide to have a biological effect endogenously, it must be resistant to degradation during intestinal transit and be able to be transported through the intestinal mucosa resisting the action of epithelial peptidases. This biological activity has a positive effect on bodily functions and conditions, and ultimately on health.

Being one of the richest sources of bioactive material, serum proteins have benefits beyond simple nutrition. These effects are, among others, antimicrobial properties, reduction of cancer cell proliferation, immune enhancement, reduction of hypertension, antioxidant and antithrombotic activity, reduction of cholesterol levels, which have beneficial effects on cancer, kidney disease, osteoporosis, obesity. Some of these bioactive peptides derived from serum proteins are lactoporphins and lactoferrins, from  $\alpha$ -LA and  $\beta$ -LG, casoplatelins, from GMP, seroporphin and albutensin A, derived from BSA, and lactoferrin and lactoferroxins from lactoferrin (Jelen, 2003).

### **3.2.1 Cottage cheese (Whey cheese)**

Whey cheese making is an ancient technique that is generally used around the world following traditional and small-scale production. It is one of the most direct forms of recovery of serum proteins, which, accompanied by certain residual fat, are denatured and coagulated after the application of heat treatment. As a general rule, if the whey is heated to more than 70°C to pH below 3.9, serum proteins precipitate and, for this

precipitation to be complete, whey must be heated to at least 90°C and kept at this temperature for several minutes (Jelen, 1992).

Whey cheeses are given different names depending on the country and region where they are produced: in Spain it is called cottage cheese; in Norway, Mysost Primost, Gjestost or Grubransdalsost; in Portugal, Requeijão; in Italy, Ricotta; in France, Serac, Brousse, Broccio, Greuil; in Greece, Manouri, Myzithra, Anthotyros; in Switzerland, Schottenziegr, Hudelziger, Mascarpone; in Israel, Urda; Ricotta being the most important and best-known cheese whey in the world, along with Mysost (Jelen, 2003).

Whey cheeses are significantly different from each other, in terms of their chemical composition, mainly due to variations in the origin and type of whey, as well as the way they are made. These cheeses are made with whey of sheep, goat and bovine origin. In Mediterranean countries they are normally based on sheep serum due to the economic importance of these ruminants in this area and their higher protein content than goat and bovine serums, which have a lower cheese yield. This type of cheese is characterized by good digestibility, being rich in protein and a high nutritional value, because it contains all the essential amino acids (especially methionine and cysteine) in sufficient quantities to meet human needs (Jelen, 2003).

Cottage cheese is mainly made from sweet whey without cream. In some cases, a small percentage of skim milk is added to the whey. Generally, it is made by heating the whey to high temperatures and adding acid, to bind the curd particles together. Whey contains the proteins lactalbumin and lactoglobulin, coagulable in heat, and in the case of skimmed milk, also casein. Precipitation takes place in such a way that the curd particles float on the surface of the whey; This is achieved by stirring and heating the serum precisely. It is a product with a soft and fragile texture (Zehren, 1976).

Technological process for the production of cottage cheese: (Zehren, 1976)

#### **a. Serum preparation**

The whey used to make cottage cheese must be sweet (not too acidic). The ideal is to use whey from the production of double cream, pear or other low-acid cheese. The acidity of the whey should be determined using the titration test. The optimal acidity of whey is 37-40°D. High acid does not result in good cottage cheese and must be neutralized with baking soda to decrease the level of acidity. The manufacture of cottage cheese is very simple to make from cheese whey from preparations that produce sweet or moderately acidic whey.

#### **b. Heating and resting of the serum**

The remaining serum is heated to approximately 85° C. This is gently shaken as the temperature rises. When the temperature reaches 85° C, about half a liter of acid whey or white vinegar is added in 5 - 6 times its volume in water per 100 liters of whey. It can also be prepared without aggregations. Acid whey should be added quickly, while vigorously shaking the contents of the tub, to prevent localized coagulation. As soon as the acidic whey has been added, the stirring stops and while the whey swirls in the vat, the curds should float on top forming a compact mass. The heating rate, the stirring method, and the acidity of the whey all influence the amount of whey to be added. Then the heating is suspended, it is left to rest until a fairly compact dough is observed.

#### **c. Drained and drained**

The albuminous clot slowly rises to the surface where it forms a thick layer that is removed by means of a skimmer or fine-mesh cloth. The whey can also be removed by siphon, opening the draining tap, being careful not to move the curd, as its particles are so fine that they are lost in the whey with movement. The curds are then placed in fine bags or canvases, which are hung or placed on a draining table and left to drain overnight (12-18 hours). It is advisable that cottage cheese drain overnight in a refrigerated room.

#### **d. Kneading and salting**

The cottage cheese is removed from the canvases or bags and kneaded perfectly. If desired, add salt to a proportion of 1 to 1.5%, knead well while mixing, to obtain a uniform dough that can be preserved for a few days.

**e. Handling cottage cheese**

After the cottage cheese is kneaded, it is packed using parchment paper or in plastic bags and sealed immediately to prevent contamination.

**f. Conservation**

Packed cottage cheese should be placed in special boxes, stored in a refrigerator at a temperature of 4 - 5° C and marketed quickly. If there is no refrigerator, it should be kept in a cool and clean place. It is a product of high humidity, approximately 80% and very perishable, so it must be consumed quickly.

### **3.2.2 Fermented milks**

The beneficial effects of consuming milk can be increased when fermented milks are consumed, especially with probiotic characteristics, that is, they have viable, non-pathogenic bacteria that exert a beneficial function in the individual. There are currently several studies, both in animals and in humans, that support the performance of the intestinal microbiota in the regulation of gastrointestinal sensory and motor function, the prevention of colorectal carcinogenesis, immunological and metabolic aspects, and behavior (Abreu, 2012), but it should be noted that the person who performs a specific action at the health level is always a specific strain and not all probiotics in general.

To produce beneficial effects on the host, probiotics do not necessarily have to colonize the target organ, although they do arrive alive in sufficient quantity to affect its microecology and metabolism. Thus, most probiotic strains are able to reach the colon alive (in a variable percentage) passing through the entire upper gastrointestinal tract, and their viability will depend on many factors: on the one hand, the intrinsic factors of the probiotic, and on the other, dependent on the host, such as, for example,

the degree of acidity in the stomach, the length of exposure to the acid, the concentration and duration of exposure to bile salts and others (Oliveira and González, 2016).

Fermented milks have proven to be nutritionally better than other dairy products such as cheese, cream, butter, etc., due to their nutritional value similar to that of milk, but with greater digestibility than milk thanks to its low pH and lower lactose concentration. In addition, in recent years there has been an increase in interest in foods with a positive effect on health beyond their nutritional value. Among them, in the field of fermented milks, probiotics stand out, partly due to the fact that this matrix is capable of keeping bacteria viable and that its consumption is recommended daily (Erdmann et al., 2008).

Fermented milks in general, and probiotics in particular, have been attributed a large number of properties, being recommended for lactose intolerance, diarrhea, constipation, presence of *Helicobacter pylori*, prevention or improvement of infections, improvement of the immune system, atopic eczema, etc. (Baró-Rodríguez et al., 2010; Khani et al., 2012; Sánchez et al., 2009). The genus *Lactobacillus* has a long history of use and safety in the manufacture of dairy products and it is for this reason that the possibly probiotic strain *Lactobacillus plantarum* C4 was chosen, isolated by our research group and which demonstrated antimicrobial and microbiota-modulating activity, as well as immunomodulatory properties (Bergillos-Meca, 2014; Bujalance et al., 2007; Fuentes et al., 2008; Puertollano et al., 2008).

During fermentation, and mainly depending on the strain used, different bioactive peptides can be released (Gobbetti et al., 2002). Biologically active peptides are peptides derived from food that exert on the body, beyond their nutritional value, a physiological effect similar to that of hormones, with fermented milks being an excellent source of them (Donkor et al., 2007; Erdmann et al., 2008). Some of these peptides have been shown to have antihypertensive, antioxidant, antibacterial, anticancer, immunomodulatory, and opioid activities, as well as mineral-

binding and regulatory capacity in metabolic syndrome (Donkor et al., 2007; Korhonen, 2009; Minervini et al., 2003; Muguerza et al., 2006; Ricci et al., 2012).

These peptides are inactive in the protein sequence, being released during gastrointestinal digestion or during food processing, such as fermentation, heating, etc. (Hernández et al., 2016). Bioactive peptides normally contain between 2 and 20 amino acids and their amino acid activity is based on their composition and amino acid sequence, on which their physicochemical properties such as charge or hydrophobicity depend (Erdmann et al., 2008). The most studied peptide activity is the inhibition of angiotensin-I converting enzyme (ACEi), which plays a crucial role in the regulation of blood pressure.

Although most of the publications on peptides with ACEi or antihypertensive activity have been made with peptides from cow's milk, in recent years goat's milk proteins are gaining importance as an alternative source of peptides with this activity (Espejo et al., 2013; Haque et al., 2007; Park et al., 2007; Ricci et al., 2012). On the other hand, some researchers have pointed out that antioxidant peptides present in food play a vital role in maintaining the body's antioxidant defenses, as well as in protecting the food from oxidation. Finally, peptides with antibacterial activity have also been discovered, useful especially for subsequent application in industry (Benkerroum, 2010).

It is considered that there are 10 bacteria in the human body for every human cell and that the intestinal microbiota fulfills three major functions: nutritive and metabolic, protective and trophic (Prados et al., 2015). Among the pathologies in which the benefits of probiotics have been demonstrated are the treatment of diarrhea, combating *Helicobacter pylori*, necrotizing enterocolitis, eliminating or reducing the effects of allergy, chronic inflammatory bowel disease and improving lipid metabolism of cholesterol and triglycerides (Vilaplana, 2015).

Probiotics help in the regulation of the immune response and their continuous use is advantageous for the health of the consumer,

participating in the modulation of the normal intestinal microbiota, reducing the risk of intestinal disorders, preventing diseases such as infections and food allergies, reducing cholesterol levels, stabilizing the intestinal mucosa and relieving the symptoms of lactose intolerance (Kich et al., 2016).

The best-known fermented milk is yogurt, a functional food obtained by fermentation of lactic acid bacteria in milk. Since ancient times, its effects on human health have been widely known, including the prevention of colon cancer, lowering cholesterol, improving intestinal microbiota, effects on the immune system, among others. The bacteria responsible for these effects are the acid-lactic-probiotic bacteria that ferment milk (Parra, 2012). Currently, there are different commercial preparations of probiotics, generally mixtures of lactobacilli and bifidobacteria (Santillán et al., 2014).

The emergence of antibiotic-resistant bacteria as well as natural ways of suppressing the growth of pathogenic microorganisms have contributed to the concept of probiotic microorganisms, which not only compete with and suppress undesirable fermentation in the human gut, and equally important produce a large number of beneficial effects on consumer health (Nagpal et al., 2012; Kailasapathy, 2013), as they act on the intestinal ecosystem, stimulating both mucosal and non-immune immune mechanisms, through antagonism and competition with potential pathogens (Villanueva, 2015).

But it is not enough for food to have enough presence of probiotic microorganisms. Consumer acceptance is a key aspect in the development of functional foods, which in general are not perceived as a separate category from natural foods, which is favorable; However, acceptance is not unconditional and the appearance and quality of the product, as well as the clarity of its statement, are important aspects in its acceptance. It has been pointed out that educational level, geographical origin and gender are variables in relation to perception and that the attitude of doctors and dieticians is important for this acceptance (Illanes, 2015).



These potential effects of fermented goat's milk could be related to the influence on adipocyte lipid metabolism and, specifically, to an increase in diet-induced thermogenesis, which would lead to higher energy expenditure and less fat storage. Another possible mechanism associated with the consumption of fermented goat's milk and the improvement of body composition could be related to the current evidence supporting the consumption of milk protein and the role of dairy in satiety, favoring weight loss and the prevention of weight gain. In this sense, fermented goat's milk induces the elevation of plasma leptin levels and the reduction of ghrelin levels, decreasing appetite, increasing the feeling of satiety and, consequently, reducing body weight (Bendtsen, 2013).

Fermentation is a simple, inexpensive, and safe way to preserve milk. In areas where modern milking and milk collection equipment is available, and where there is generally a great deal of knowledge and experience in raw milk preservation techniques, and where they also have good transport and distribution systems, there is no need to use fermentation as a preservation method. Conversely, in areas or countries that do not have all of these mediums, milk fermentation as a means of preservation still retains the importance it originally had. Lactic acid bacteria change the characteristics of milk, so that most undesirable microorganisms, including pathogens, cannot grow in it, or even die.

Among the changes that occur in milk is the decrease in pH (up to 4.6-4), a factor that contributes to the maintenance of a low pH in the stomach after consuming milk; inhibition of microbial development by undissociated acids (e.g., lactic acid), and by other metabolites such as H<sub>2</sub>O<sub>2</sub> and other substances with antibiotic activity; low oxide-reduction potential; and the consumption by lactic acid bacteria of components that are vital for other microorganisms.

Proper pasteurization of raw milk destroys any pathogens that might survive fermentation. There are many different fermented milks, but when it comes to the technology of their manufacture, they are all similar. Fermented milks can be classified in several ways, but their classification

is generally accepted according to the type of microorganism used in their production (Pérez and Sánchez, 2012).

## Chapter IV

### Technology for the production of fermented milks

The technology of fermented milks is simple and small-scale brewing requires only remarkably simple equipment. Large-scale manufacturing requires consistent, low-cost production, which requires greater control and more sophisticated equipment, although the basic manufacturing principles are the same. There are many types of fermented milk that are made using similar technology and, in many cases, the differences are limited to the type of starter culture and the total solids content of the milk (Pérez and Sánchez, 2012).

Traditionally, yogurt is made using the starter cultures *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus*. These bacteria provide man with some health benefits, but they are not natural inhabitants of the intestinal tract and do not survive under high concentrations of acid and bile salts. The solution has been to incorporate probiotic bacteria into fermented dairy products because these are live microbial cultures, used as dietary supplements or food ingredients which, when ingested, produce a beneficial health effect to the host, improving the balance of the flora of the gastrointestinal tract, inhibiting or preventing the growth of pathogenic bacteria that cause diseases (Hernández et al., 2016).

The general principle of processing methods in goat's milk are the same as those used in cow's milk, which consist of reducing the pH and activity of water to prolong its shelf life. The acid gel of goat's milk is characterized by firmness and lower viscosity compared to cow's and sheep's milk.

The viscosity of the gel is associated with the casein content in milk, especially with the  $\alpha$ -s fraction casein, which is in goat's milk from 25 to 30%. In addition, the appropriate heat treatment, the addition of

stabilizers and the type of mother culture applied, are the other factors that reduce the syneresis intensity (Hernández et al., 2016).

## 4.1 Types of fermented milks

### 4.1.1 Fermented milks with thermophilic lactic acid bacteria

They are the most commercialized products. The microorganisms responsible are strains of *Lactobacillus delbrueckii* subsp. *bulgaricus*, thermophilic bacteria whose optimal growth temperature is between 42 - 43°C. During lactic fermentation carried out by these bacteria, metabolites such as acetaldehyde and diacetyl are produced that give them a characteristic taste and aroma. Lactic acid is also produced until pH values of 3.8 – 4 are reached. This increase in acidity causes the casein in milk to coagulate, improving its preservation (Sampablo, 2019).

- a. **Acidophilic milk:** Originally from the USA. It is milk fermented by *Lactobacillus acidophilus* with a curdled, mixed or liquid texture and a mild flavor.
- b. **Yoghurt or yoghourt:** A coagulated milk product obtained by lactic fermentation by the action of *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* from milk or concentrated milk, whether or not skimmed, or from cream, or from a mixture of two or more of these products, with or without the addition of other milk ingredients, which have previously undergone heat treatment or other treatment, equivalent, at least, to pasteurization. All the microorganisms producing lactic fermentation must be viable and present in the milk part of the finished product in a minimum quantity of 10<sup>7</sup> colony-forming units (CFU) per gram or millilitre.

Depending on the products to be added, or the application of heat treatment after fermentation, different types of yogurts appear: sweetened yogurt, sweetened yogurt, flavored yogurt, yogurt with fruit, juice and/or other foods, pasteurized yogurt after fermentation. Pasteurized yoghurt after fermentation meets all the requirements established in the definition of yoghurt except for the viability of the microorganisms producing

fermentation, as it has been subjected to a subsequent thermal process by which the lactic acid bacteria it contained have lost their viability (Sampablo, 2019).

#### **4.1.2 Fermented milks with mesophilic lactic acid bacteria**

- a. **Filmjöl**: It originates in Sweden but is consumed in northern Europe in general. The bacteria involved in fermentation are mesophilic, i.e. their optimal growth temperature is between 20-22°C. Among them *Lactococcus lactis*, *Lactococcus lactis cremoris*, *Lactococcus lactis diacetylactis* and *Leuconostoc mesenteroides cremoris*.

#### **4.1.3 Fermented milks with lactic acid bacteria and yeasts**

They are produced by the heterolactic fermentation of milk by the action of bacteria (lactic fermentation) and yeasts (alcoholic fermentation). They are slightly alcoholic beverages (up to 2% ethanol), sparkling, due to the CO<sub>2</sub> produced and acidic as a result of the lactic acid generated.

- a. **Kefir**: Its alcohol content is low (in order of 0.5%). Bacteria of the genera *Lactobacillus*, *Leuconostoc* and *Lactococcus* and yeasts are involved in the fermentation of milk; both lactose fermenters (*Kluyveromyces marxianus*) and yeasts that ferment without the need for lactose (*Saccharomyces sp.*) (Sampablo, 2019).

- b. **Kumys**: It is native to Mongolia. Traditionally it was made with mare's milk, but today it is made with cow's milk. It contains more alcohol (up to 3%) than kefir, due to sucrose's addition to milk. Bacteria such as *Lactobacillus acidophilus* and *Lactobacillus delbrueckii subsp. bulgaricus* and yeasts such as *Kluyveromyces marxianus* are involved in fermentation (Sampablo, 2019).

#### **4.1.4 Fermented milks with lactic acid bacteria and moulds**

- a. **Villi**: is a viscous fermented milk, native to Finland. The starter culture is made up of lactic acid bacteria and molds such as *Geotrichum candidum*. The milk used as raw material is not homogenized after

pasteurization, so the cream is separated on the surface and is where mold develops. During fermentation, this mold produces a large amount of CO<sub>2</sub> (Sampablo, 2019).

## b. Condensed milk

Condensed milk Condensed milk is a by-product derived from whole milk, which is obtained by partially evaporating the water in the milk with the addition of sugar, helping to preserve it. Due to its high energy content (Table 1), condensed milk has been used in the food industry as a sweetener for fruits, condensed milk, due to its high sucrose content, inactivates the microorganisms present in the product, which is not highly recommended as a basic food for healthy diets. In the nutritional field, it is not highly recommended to consume it in high quantities, due to the high sugar content it contains, in this case children could have dental problems and in the future it could cause chronic diseases such as obesity, diabetes (Gavino, 2021).

**Table 1. Composition of condensed milk**

Nutrients	Unit	Value per 1000 grams
Water	g	27.16
Energy	kcal	321
Proteins	g	7.91
Total lipids	g	8.7
Carbohydrates	g	54.4
Total sugars	g	54.4
<b>Minerals</b>		
Calcium	mg	284
Iron	mg	0.19

Magnesium	mg	26
Phosphorus	mg	253
Potassium	mg	371
Sodium	mg	127
Zinc	mg	0.94
<b>Vitamins</b>		
Vitamin C	mg	2.6
Thiamine	mg	0.09
Riboflavin	mg	0.416
Niacin	mg	0.21
B6	mg	0.051
Folate	µg	11
B12	µg	0.44
A	µg	74
And	µg	0.16
D	µg	0.2
K	µg	0.6
<b>Lipids</b>		
Total saturated fatty acids	g	5.486
Total monounsaturated fatty acids	g	2.427
Total polyunsaturated fatty acids	g	0.337
Cholesterol	mg	34

Source: (Gavino, 2021)

## 4.2 Condensed milks

- a. Crystallization Lactose in milk is present in two isomeric forms which are called  $\alpha$  and  $\beta$ lactose, the  $\alpha$  form tends to crystallize due to its low solubility which tends to crystallize in products such as condensed milk this is due to the solubility of lactose decreases due to the high sugar content that is added in condensed milks, which can lead to the crystallization of condensed milks (Verhelst, 2015).
- b. Thickening The main physical defect of condensed milk is the thickening or change in viscosity, this is due to transitions in the composition of the milk or its heat treatment by the addition of sugar or stabilizing salts (Verhelst, 2015).
- c. Color change The color change due to the effect of its heat treatment at the time of pasteurization, the presence of the golden color increases because the pH increases above 7.0, which can be solved with a proper heat treatment in the proper evaporation of the milk. The appearance of color is due to the evaporation time and the volume of the mixture and the heat treatment that is carried out manually or with appropriate equipment (Verhelst, 2015).

### 4.2.1 Technological process for the production of condensed milk

According to (López, 2022), the following procedure must be carried out for the production of condensed milk:

- Reception

Reception is carried out in storage tanks.

- Filtration

In this part of the process, a cloth or membrane is used to filter the milk and all impurities present at the time of milking the goat are removed.

- Pasteurization

Goat's milk is heated to a temperature of approximately 65°C, in this process all the microorganisms present in the milk are inactivated.



- Addition

Sugar is added by making a separate dilution for better homogenization, mixing it homogeneously to avoid generating lumps in the pot at a temperature of 70°C for about 60 min.

- Concentration

The temperature is lowered to approximately 50°C until it acquires the required texture, this stage is the important one to consider the temperature and homogenization because if it is at a high temperature it tends to burn.

- Cooling

The temperature of the condensed goat's milk is lowered to 15°C for 40 to 60 min.

- Packaging

The finished product is packaged in heat-resistant glass jars which were sterilized before finishing the process of making condensed goat's milk to avoid contamination when in contact with it.

#### **4.2.2 Butter**

Butter can be defined as cream or simply as high-fat milk. The process of obtaining the cream is carried out by simple means such as keeping the whole milk at rest for a certain period of time in order to ensure that the fat rises due to the effects of gravity. The fat of the milk has a lower density than the rest of the components of the milk and for this characteristic it rises to the surface of the milk forming what is called (the cream). Under these conditions, this element can be removed by means of ladles and the resulting product is called cream. Another way to obtain cream, already at an industrial level, is by using centrifugal force.

In fact, equipment is manufactured equipped with rotating plates with special characteristics that are intended to separate the heavier elements of milk from the lighter ones. In this way, the fat is separated from the rest

of the components of the milk. The resulting product, as in the previous case, is called cream. There are two types of cream depending on the degree of acidity it has and consequently, the type of butter you want to obtain.

Thus, we speak of sweet cream and sour cream. Sweet cream is one whose acidity remains at levels similar to those of normal milk. On the other hand, sour cream is one that, as a result of the fermentation of lactose by the action of microorganisms, has increased levels of acidity. And, at the same time, the taste and pleasant smell of it has been enhanced, a phenomenon that is transmitted to butter (Guzmán, 2000).

Butter obtained from sour creams, although it is true that it is better accepted by the consumer, has a shorter shelf life. The reason for this phenomenon is that the product can reach what is called over-ripening due to the action of microorganisms incorporated during the inoculation and maturation process. For this reason, companies choose to produce butters from pasteurized cream to which lactic ferments have not been incorporated and it is thus that in the market it is difficult to find the product obtained from acid creams (Guzmán, 2000).

#### **4.2.3 Stages of production of goat's milk butter**

It corresponds to the reception and quality control of the raw material. The main controls to be carried out at this stage are the determination of the acidity of the cream and especially the temperature control. The process of making butter requires the use of temperatures that do not exceed 80C.

- The cream cooled to 8°C undergoes the whipping process. The specifications for this stage are detailed above.
- Once the beating stage is finished, the grain is washed. At least three washes are carried out. The last wash is done with salt water. The amount of salt corresponds to 3% of 10 kilos of butter calculated.
- Kneading, the stage after washing, has among its objectives to distribute salt, regulate humidity and improve the texture of the

product. The butter is kneaded until the humidity is not higher than 16%.

- Once the kneading is finished, the butter is molded, packaged and stored. The storage temperature should be no higher than 50C.
- Unconventional derivatives of goat's milk are lesser-known but equally valuable dairy products, such as fermented milks, whey, cottage cheese, condensed milk, and butter. These products have a unique taste and texture that make them popular with consumers looking for different and authentic dairy products.
- Unconventional goat's milk derivatives are also an excellent source of nutrients, such as protein, calcium, and vitamins, making them a healthy alternative to conventional dairy products.
- The production of these products requires a high level of skill and knowledge to ensure the quality and safety of the final product, especially in the case of fermented milks and butter.
- Despite their lower awareness and marketing, unconventional derivatives of goat's milk are gaining popularity due to their distinctive taste and nutritional benefits, which has led to an increase in the production and marketing of these products worldwide.

In theory, unconventional goat's milk derivatives are valuable and unique dairy products that offer a healthy and authentic alternative to conventional dairy products. Its production requires skill and knowledge to ensure the quality and safety of the final product, but its popularity is on the rise due to its distinctive flavor and nutritional benefits.

### **4.3 Navigating the Regulatory Landscape and Ensuring Quality Assurance in Goat Milk Development**

Goat milk has emerged as a significant alternative to cow's milk, gaining popularity among consumers for its unique nutritional profile, digestibility, and lower lactose content. Rich in essential fatty acids, vitamins, and minerals, goat milk is not only suitable for individuals with

lactose intolerance and equally important offers numerous health benefits recognized across various cultures for centuries.

The increasing awareness of health and wellness, combined with rising dietary preferences such as veganism and dairy-free lifestyles, has paved the way for goat milk to establish itself in the dairy market. This development is not merely a trend; it represents a shift in consumer attitudes toward diverse sources of nutrition and sustainable farming practices. As more people seek alternatives that align with their health goals and ethical beliefs, the demand for goat milk products continues to rise.

However, the growth of the goat milk industry is accompanied by a complex regulatory landscape that governs its production, processing, and distribution. Understanding the various regulations—both at national and international levels—is essential for stakeholders in the goat milk market, from farmers to processors and retailers. Compliance with these regulations ensures the safety, quality, and consistency of goat milk products, which are vital for building consumer trust and achieving market success.

Also, quality assurance measures are paramount in the processing of goat milk. Implementing rigorous hygiene practices, testing protocols, and certification programs not only help meet regulatory requirements on the contrary enhance product quality, ensuring that consumers receive safe and nutritious options.

As we delve into the intricacies of the regulatory landscape and quality assurance measures necessary for goat milk development, it is crucial to recognize the challenges and opportunities that lie ahead. Understanding these elements will provide insights into how the goat milk industry can thrive in a competitive market while addressing the growing demands of consumers for transparency, sustainability, and quality.

#### **a. Regulatory Landscape for Goat Milk Production**

The regulatory landscape for goat milk production is a crucial framework that ensures the safety, quality, and consistency of goat milk products. Given the increasing popularity of goat milk as an alternative to cow's milk, understanding the various regulations at national, international, state, and local levels is essential for producers, processors, and consumers alike.

### **b. Overview of National Regulations**

In the United States, goat milk production is primarily governed by the Food and Drug Administration (FDA) and the United States Department of Agriculture (USDA). The FDA sets the standards for the pasteurization and labeling of milk and milk products, including goat milk. According to the FDA's Pasteurized Milk Ordinance (PMO), all goat milk intended for sale must be pasteurized to eliminate pathogens that could pose health risks (U.S. Department of Health and Human Services, 2019). Additionally, the FDA requires that goat milk products adhere to strict labeling guidelines, which include accurate ingredient lists and nutritional information to ensure consumer safety and informed choices.

State regulations also play a significant role in goat milk production. Each state has its own agricultural department that may impose additional standards beyond federal guidelines. These regulations can cover everything from herd health management to facility hygiene. Producers must navigate these state-specific requirements, which can vary widely, to ensure compliance and maintain their ability to market their products.

#### **4.3.1 International Standards and Compliance**

As goat milk gains traction in global markets, international standards for food safety and quality assurance have become increasingly relevant. Organizations such as the Codex Alimentarius Commission, established by the World Health Organization (WHO) and the Food and Agriculture Organization (FAO), provide a set of internationally recognized standards that govern the production, processing, and trade of food products,

including goat milk (Moises et al., 2024). Compliance with these standards is essential for producers aiming to export their products or compete in international markets.

International standards typically encompass guidelines related to hygiene, production practices, and labeling. Adhering to these standards not only helps ensure product safety and equally important facilitates smoother trade by aligning with the requirements of importing countries. As countries increasingly demand high-quality and safe food products, understanding and complying with these international regulations becomes imperative for goat milk producers.

In addition to federal and international regulations, state and local laws significantly impact goat milk production. Local health departments often establish specific sanitation and health codes that must be followed by producers and processors. These regulations may dictate everything from the construction of milking facilities to the management of waste byproducts.

Some states have enacted specific legislation aimed at promoting the dairy goat industry. To illustrate, programs may be in place to support research and development, provide financial assistance for small-scale farmers, or encourage the use of sustainable farming practices. Understanding and adapting to these state and local regulations is crucial for goat milk producers seeking to thrive in a competitive market.

Indeed, the regulatory landscape for goat milk production is multifaceted, encompassing national, international, state, and local regulations. Producers must remain informed and compliant with these diverse regulations to ensure product safety and quality while effectively navigating the complexities of the goat milk market.

#### **4.3.2 Quality Assurance Measures in Goat Milk Processing**

In the realm of goat milk development, quality assurance is paramount. It not only safeguards consumer health but also enhances the marketability of goat milk products. Effective quality assurance measures

are essential in ensuring that goat milk is safe, nutritious, and free from contaminants. This section delves into the critical components of quality assurance in goat milk processing, focusing on hygiene and sanitation practices, testing and quality control procedures, and the significance of certification programs.

The foundation of quality assurance in goat milk processing begins with stringent hygiene and sanitation practices. Maintaining a clean environment is crucial for preventing the introduction of pathogens and contaminants into the milk supply. This involves regular cleaning and sanitization of equipment, milking facilities, and storage areas.

Personnel involved in goat milk production must adhere to strict hygiene protocols, including handwashing and wearing appropriate protective gear. Therefore, the goats themselves should be kept in clean, well-ventilated environments to reduce the risk of disease transmission and ensure the production of high-quality milk (Ulsenheimer et al., 2022). Implementing a comprehensive sanitation schedule, which includes routine inspections and maintenance of equipment, is vital for sustaining hygiene standards throughout the processing chain.

Testing and quality control procedures are integral to ensuring the safety and quality of goat milk products. Regular testing for microbiological contaminants, such as bacteria and viruses, is essential to verify that milk meets safety standards. Chemical analyses are also conducted to check for residues of antibiotics, pesticides, and other harmful substances that could compromise product integrity.

Quality control procedures should encompass a variety of parameters, including fat content, protein levels, and overall sensory characteristics. By employing standardized testing methods, producers can ensure that their products consistently meet regulatory requirements and consumer expectations. Furthermore, implementing a robust traceability system allows producers to monitor the journey of goat milk from farm to consumer, enabling swift action in the event of quality concerns.

Certification programs play a crucial role in the goat milk industry by establishing standards for quality and safety. These programs, which may be national or international in scope, provide frameworks for producers to demonstrate compliance with best practices in goat milk processing. Common certifications include organic, non-GMO, and animal welfare certifications, which can enhance consumer trust and expand market access.

Participating in certification programs not only helps producers meet regulatory requirements but also positions them favorably in an increasingly competitive marketplace. As consumers become more discerning about the products they purchase, certifications serve as valuable indicators of quality and ethical practices. Furthermore, engaging in these programs can drive continuous improvement, encouraging producers to adopt innovative practices and technologies that enhance product quality and safety.

In summary, quality assurance measures in goat milk processing are multifaceted and essential for the success of the industry. By prioritizing hygiene and sanitation, implementing rigorous testing and quality control procedures, and participating in certification programs, producers can ensure that they deliver safe, high-quality goat milk products to consumers. As the demand for goat milk continues to grow, maintaining these standards will be critical for sustaining consumer confidence and fostering industry growth.

#### **4.4 Future Directions in Goat Milk Development**

The goat milk industry has experienced significant growth over the past few decades, driven by increasing consumer awareness of its nutritional benefits and versatility. However, several challenges persist that could hinder further development. Addressing these challenges while capitalizing on emerging opportunities will be essential for the sustainable advancement of goat milk production and marketing.



One of the primary challenges facing the goat milk industry is fluctuating market demand. While interest in goat milk has risen, driven by trends favoring alternative dairy products, competition from cow's milk and other plant-based beverages remains strong. Consumer preferences are constantly evolving, influenced by health trends, dietary restrictions, and ethical considerations (Miller and Lu, 2019). Therefore, producers must stay attuned to these trends and adapt their marketing strategies to effectively communicate the unique benefits of goat milk, such as its digestibility and lower allergenic potential compared to cow's milk.

Additionally, educating consumers about the nutritional advantages of goat milk—like its rich content of vitamins, minerals, and beneficial fatty acids—can help increase its appeal. Innovative marketing approaches, including collaborations with health-focused influencers and participation in farmer's markets, could enhance visibility and foster a loyal customer base.

To meet rising demand and improve production efficiency, the goat milk industry must embrace innovation. Advances in breeding techniques, animal husbandry practices, and nutrition can lead to higher milk yields and better quality products. Genetic selection for traits such as disease resistance and milk composition can enhance the productivity and sustainability of goat herds.

But, leveraging technology in goat milk processing—such as automated milking systems and state-of-the-art pasteurization methods—can improve product safety and consistency. Research into alternative processing techniques, such as fermentation and fortification, can also expand the product range and cater to diverse consumer preferences.

Sustainability is an increasingly critical concern in all areas of agriculture, including goat milk production. As consumers become more environmentally conscious, they are more likely to support brands that demonstrate responsible practices. Goat farming can offer several

environmental benefits, such as lower greenhouse gas emissions compared to cattle farming and the ability to thrive on marginal lands unsuitable for other livestock.

However, producers must also address challenges related to land use, waste management, and water consumption. Implementing sustainable practices, like rotational grazing and organic farming methods, can help mitigate negative environmental impacts and enhance the overall sustainability of goat milk production.

In synthesis, while the goat milk industry faces several challenges—including fluctuating market demands and environmental sustainability—it also has numerous opportunities for growth through innovation and consumer education. By addressing these issues head-on, the industry can pave the way for a more prosperous and sustainable future for goat milk development.

As we look toward the future of goat milk development, it is evident that this sector is poised for significant growth driven by evolving consumer preferences, increasing awareness of health benefits, and innovative production techniques. The regulatory landscape, while complex, continues to adapt in response to these trends, ensuring that safety and quality remain paramount in the production of goat milk products (Moises et al., 2024).

As more consumers seek lactose-free options and are drawn to the perceived health advantages of goat milk, producers are presented with unique opportunities to expand their market reach. This surge in demand necessitates a focus on not only meeting regulatory standards but also enhancing the overall quality and safety of products.

Innovations in production techniques—including advancements in breeding, feed optimization, and sustainable farming practices—are helping to improve yield and efficiency while maintaining ethical standards. These innovations are critical as they align with consumer

expectations for transparency and sustainability, which are increasingly influencing purchasing decisions.

Moreover, the emphasis on quality assurance measures — ranging from stringent hygiene practices to rigorous testing protocols — will continue to play a crucial role in ensuring that goat milk products meet the highest standards of safety and quality. Certification programs will further bolster consumer confidence, allowing producers to differentiate their products in a competitive marketplace.

The future development of goat milk is bright, marked by an intersection of regulatory compliance, quality assurance, and consumer-driven innovation. As industry navigates challenges and embraces opportunities, stakeholders must remain committed to sustainable practices that not only enhance product quality on the contrary contribute positively to environmental stewardship. With a proactive approach to regulation and a focus on quality, the goat milk industry is well-positioned to thrive in the years ahead, meeting the demands of an increasingly health-conscious and environmentally aware consumer base.

## Conclusion

Goat milk offers unique flavor and potential health benefits, making it a noteworthy alternative to cow milk. Key advantages include:

- *Allergy-Friendly*: Its different protein structure may cause fewer allergic reactions, making it suitable for those allergic to cow milk.
- *Lactose Intolerance*: Goat milk has lower lactose content and smaller fat globules, aiding digestion for many lactose-intolerant individuals, though caution is advised for those with severe intolerance.
- *Nutritional Value*: Rich in essential nutrients like calcium, phosphorus, and vitamins A and D, goat milk is particularly beneficial for children and the elderly. Its digestibility supports nutrient absorption, essential for older adults.

In culinary applications, goat milk is versatile, used in cheeses, yogurt, and desserts, and is popular in Mediterranean and Middle Eastern cuisines. The trend towards sustainable dairy sources has increased its demand. Despite its benefits, individual dietary needs and potential allergies should be considered. Consulting healthcare professionals is advisable for significant dietary changes. Comparatively, goat milk has higher protein and superior digestibility than cow milk, making it an excellent alternative. It also boasts a robust nutrient profile with higher calcium and certain B vitamins. Other milks, like sheep and camel, offer unique advantages, but goat milk stands out for its versatility and nutrition.

Goat milk is a rich source of essential vitamins and minerals like calcium, magnesium, phosphorus, and potassium, vital for bone health, cardiovascular function, and muscle contraction. It also contains significant vitamins such as riboflavin (B2), vitamin B12, and vitamin D, important for energy metabolism, red blood cell formation, and calcium absorption. The presence of beneficial fats enhances the bioavailability of these nutrients. This nutrient profile supports the potential for creating various

unconventional derivatives, including fortified foods and dietary supplements. Goat milk's unique biochemical compounds further position it as a valuable resource for innovative applications.

Goat milk is valued for its unique biochemical compounds, enhancing its versatility in various applications:

- *Short-Chain Fatty Acids (SCFAs)*: Goat milk has a higher concentration of SCFAs, like butyric and caproic acid, which support gut health and provide rapid energy. They also enhance flavor, making goat milk ideal for specialty culinary products.

- *Bioactive Peptides*: Produced during digestion, these peptides offer health benefits like antihypertensive and antimicrobial effects. The unique protein composition of goat milk enhances the bioavailability of these peptides, making it suitable for dietary supplements and fortified foods.

- *Antioxidants and Immunoglobulins*: Goat milk is rich in antioxidants (selenium, vitamin E) and immunoglobulins (IgG), supporting immune function and combating oxidative stress. These properties make it appealing for health-focused consumers and functional food development.

In summary, the distinctive compounds in goat milk not only improve its nutritional profile but also open opportunities for innovative applications in food and health industries. Indeed, the regulatory landscape for goat milk production is multifaceted, encompassing national, international, state, and local regulations. Producers must remain informed and compliant with these diverse regulations to ensure product safety and quality while effectively navigating the complexities of the goat milk market.

In theory, unconventional goat's milk derivatives are valuable and unique dairy products that offer a healthy and authentic alternative to conventional dairy products. Its production requires skill and knowledge to ensure the quality and safety of the final product, but its popularity is on the rise due to its distinctive flavor and nutritional benefits. As we consider the future, the dairy sector is undergoing significant changes,

adjusting to new challenges and opportunities. The adoption of cutting-edge technologies, including automation, artificial intelligence, and data analytics, is transforming production methods in goat milk, improving quality assurance, and streamlining supply chain operations.

Furthermore, the increasing focus on sustainability is driving the industry to investigate environmentally friendly practices, such as minimizing waste and implementing energy-efficient processing techniques. These developments not only address consumer demands but also align with worldwide initiatives aimed at combating climate change and fostering environmental responsibility.

## Bibliography

- Abreu, A. T. (2012). Prebióticos, probióticos y simbióticos. *Revista de gastroenterología de Mexico*, 77, 26-28.
- Aider, M., Halleux, D., Melnikova, I. (2009). Skim acidic milk whey cryconcentration and assessment of its functional properties: Impact of processing conditions. *Innovative Food Science and Emerging Technologies*, 10(3), 334- 341.
- ALKaisy, Q. H., Al-Saadi, J. S., Al-Rikabi, A. K. J., Altemimi, A. B., Hesarinejad, M. A., & Abedelmaksoud, T. G. (2023). Exploring the health benefits and functional properties of goat milk proteins. *Food science & nutrition*, 11(10), 5641–5656.  
<https://doi.org/10.1002/fsn3.3531>
- Almasri, R. S., Bedir, A. S., Ranneh, Y. K., El-Tarabily, K. A., & Al Raish, S. M. (2024). Benefits of Camel Milk over Cow and Goat Milk for Infant and Adult Health in Fighting Chronic Diseases: A Review. *Nutrients*, 16(22), 3848. <https://doi.org/10.3390/nu16223848>
- Araujo, H. C. S., de Jesus, M. S., Sandes, R. D. D., Leite Neta, M. T. S., & Narain, N. (2024). Functional Cheeses: Updates on Probiotic Preservation Methods. *Fermentation*, 10(1), 8.  
<https://doi.org/10.3390/fermentation10010008>
- Ballard, O., & Morrow, A. L. (2013). Human milk composition: nutrients and bioactive factors. *Pediatric Clinics of North America*, 60(1), 49-74.
- Baró Rodríguez, L., Lara Villoslada, F., & Corral Román, E. (2010). *Capítulo 1. Leche y derivados lácteos*. In Á. Gil & M. D. Ruiz-López (Eds.), *Tratado de Nutrición. Tomo II. Composición y calidad nutritiva de los alimentos* (Second., pp. 1–26). Madrid: Editorial Médica Panamericana.
- Bendtsen, L. Q., Lorenzen, J. K., Bendtsen, N. T., Rasmussen, C., & Astrup, A. (2013). Effect of dairy proteins on appetite, energy expenditure,

body weight, and composition: a review of the evidence from controlled clinical trials. *Advances in nutrition (Bethesda, Md.)*, 4(4), 418–438. <https://doi.org/10.3945/an.113.003723>

- Benkerroum, N. (2010). Antimicrobial peptides generated from milk proteins: a survey and prospects for application in the food industry. A review. *International Journal of Dairy Technology*, 63(3), 320-338.
- Bergillos-Meca, T. (2014). *Desarrollo de una leche fermentada con una cepa probiótica de Lactobacillus plantarum: biodisponibilidad mineral y efectos sobre la barrera de permeabilidad intestinal y la microbiota fecal* [Tesis de doctorado]. Universidad de Granada.
- Bujalance, C., Moreno, E., Jimenez-Valera, M., & Ruiz-Bravo, a. (2007). A probiotic strain of *Lactobacillus plantarum* stimulates lymphocyte responses in immunologically intact and immunocompromised mice. *International Journal of Food Microbiology*, 113(1), 28-34.
- Caro, I., Soto, S., Franco, M. J., Meza, M., Alfaro, R. H., Mateo, J. (2011). Composition, yield, and functionality of reduced-fat Oaxaca cheese: Effects of using skim milk or a dry milk protein concentrate. *J. Dairy Sci.*, 94, 580-588.
- Donkor, O. N., Henriksson, A., Singh, T. K., Vasiljevic, T., & Shah, N. P. (2007). ACE-inhibitory activity of probiotic yoghurt. *International Dairy Journal*, 17(11), 1321-1331.
- Erdmann, K., Cheung, B. W. Y., & Schröder, H. (2008). The possible roles of food-derived bioactive peptides in reducing the risk of cardiovascular disease. *The Journal of Nutritional Biochemistry*, 19(10), 643-654.
- Espejo-Carpio, F. J., Pérez-Gálvez, R., Guadix, E. M., & Guadix, A. (2013). Optimisation of the hydrolysis of goat milk protein for the



production of ACE-inhibitory peptides. *The Journal of Dairy Research*, 80(2), 214-222.

- Facioni, M. S., Raspini, B., Pivari, F., Dogliotti, E., & Cena, H. (2020). Nutritional management of lactose intolerance: the importance of diet and food labelling. *Journal of translational medicine*, 18(1), 260. <https://doi.org/10.1186/s12967-020-02429-2>
- FAO. (2009). Food and Agriculture Organization of the United Nations.
- FAO. (1980). Manual de elaboración de quesos. Equipo Regional de Fomento y Capacitación en Lechería para América Latina. Valdivia, Chile.
- Fuentes, S., Egert, M., Jimenez-Valera, M., Monteoliva-Sanchez, M., Ruiz-Bravo, A., & Smidt, H. (2008). A strain of *Lactobacillus plantarum* affects segmented filamentous bacteria in the intestine of immunosuppressed mice. *FEMS Microbiology Ecology*, 63(1), 65-72.
- Fernández, M., Fornari, M., Mazutti, D., Olivera, F., Ferreira, A., Cichoski, R., Cansian, M., Luccio., Treichel, H. (2009). Production and characterization of xantham gum by *xanthomonas campestris* usin cheese whey as sole carbon source. *Journal of Food Engineering*, 50(2), 119-123.
- Gavino Ospino, E. O. (2021). *Uso de quimosina en el desarrollo de un método rápido de elaboración de leche condensada* [Tesis de Maestría]. Universidad Nacional José Faustino Sánchez Carrión.
- Gobbetti, M., Stepaniak, L., De Angelis, M., Corsetti, a, & Di Cagno, R. (2002). Latent bioactive peptides in milk proteins: proteolytic activation and significance in dairy processing. *Critical Reviews in Food Science and Nutrition*, 42(3), 223-239.
- Golightly, L., Chang, V., Xu, Q.A., Gao, X., & Liu, B.S. (2022). Adoption of cloud computing as innovation in the organization. *International*

*Journal of Engineering Business Management*, 14.

<https://doi.org/10.1177/18479790221093992>

- González, M. (2010). The biotechnological utilization of cheese whey. *Bioresource Technology*, 57, 11-73.
- Guzmán, V. (2000). Elaboración de productos con leche de cabra. Fundación para la Innovación Agraria Ministerio de Agricultura, Santiago de Chile.
- Haque, E., & Chand, R. (2007). Antihypertensive and antimicrobial bioactive peptides from milk proteins. *European Food Research and Technology*, 227(1), 7-15.
- Hernández, A., Torres, A., Duarte, C., Rodríguez, D. (2016). Desarrollo de una leche fermentada de cabra con cultivos probióticos. *Tecnología Química*, 36(3), 321-335. Available at <https://www.redalyc.org/pdf/4455/445546669002.pdf>.
- Holsinger, V.H., Posati, L.P., & DeVilbiss, E.D. (1974). Whey Beverages. *Journal of Dairy Science*, 57(8), 849-859.
- Huang, X. (2024). The Promotion of Anaerobic Digestion Technology Upgrades in Waste Stream Treatment Plants for Circular Economy in the Context of “Dual Carbon”: Global Status, Development Trend, and Future Challenges. *Water*, 16(24), 3718. <https://doi.org/10.3390/w16243718>
- Illanes, A. (2015). Alimentos funcionales y biotecnología. *Revista Colombiana de Biotecnología*, 17(1), 5-8.
- Jelen, P. (1992). Whey: composition, properties, processing and uses. En Y.H. Hui, *Encyclopedia of food science and technology*, 2835-2845.
- Jelen, P. (2003). Whey processing. Utilization and Products..*Encyclopedia of Dairy Science*, 25(3), 105-126.

- Kailasapathy, K. (2013). Commercial sources of probiotic strains and their validated and potential health benefits-a review. *International Journal of Fermented Foods*, 2(1), 1.
- Kalogiannidis, S., Kalfas, D., Papaevangelou, O., Giannarakis, G., & Chatzitheodoridis, F. (2024). The Role of Artificial Intelligence Technology in Predictive Risk Assessment for Business Continuity: A Case Study of Greece. *Risks*, 12(2), 19.  
<https://doi.org/10.3390/risks12020019>
- Khani, S., Hosseini, H., Taheri, M., Nourani, M., & Imani Fooladi, A. (2012). Probiotics as an alternative strategy for prevention and treatment of human diseases: a review. *Inflamm Allergy Drug TArgets*, 11, 79–89.
- Korhonen, H. (2009). Milk-derived bioactive peptides: From science to applications. *Journal of Functional Foods*, 1(2), 177–187.
- Kreczmann, B., Alonso, A., Liloia, M., Zamboni, E., Cerutti, R., Baroni, D., y Poluján D. (2015) Procesamiento del lactosuero: elaboración de lactosa y aprovechamiento de proteínas. *Procesos Tecnología Láctea Latinoamericana*. 87. Available at  
[https://www.academia.edu/checkout?feature=ALL\\_READER\\_PRODUCTS&trigger=swp-sticky-header-work-card-download-button&upgradeRedirect=%2F%2Fwww.academia.edu%2F37937970%3FbulkDownload%3Dtrue](https://www.academia.edu/checkout?feature=ALL_READER_PRODUCTS&trigger=swp-sticky-header-work-card-download-button&upgradeRedirect=%2F%2Fwww.academia.edu%2F37937970%3FbulkDownload%3Dtrue)
- Kich, D. M., Vincenzi, A., Majolo, F., de Souza, C. F. V., & Goettert, M. I. (2016). Probiotic: effectiveness nutrition in cancer treatment and prevention. *Nutrición Hospitalaria*, 33(6), 1430-1437.
- López, L. E. (2022). *Formulación de leche condensada de cabra (capra aegrus hircus) con harinas de amaranto (amaranthus spp.), quinua (chenopodium quinoa) y arroz (oryza sativa)* [Tesis de grado]. Universidad Agraria del Ecuador.

- Magableh, K. N. Y., Kannan, S., & Hmoud, A. Y. R. (2024). Innovation Business Model: Adoption of Blockchain Technology and Big Data Analytics. *Sustainability*, 16(14), 5921.  
<https://doi.org/10.3390/su16145921>
- Miller, B. A., & Lu, C. D. (2019). Current status of global dairy goat production: an overview. *Asian-Australasian journal of animal sciences*, 32(8), 1219–1232. <https://doi.org/10.5713/ajas.19.0253>
- Minervini, F., Algaron, F., Rizzello, C. G., Fox, P. F., Monnet, V., & Gobbetti, M. (2003). Angiotensin I-Converting-Enzyme-Inhibitory and Antibacterial Peptides from Lactobacillus Caseins of Milk from Six Species Angiotensin I-Converting-Enzyme-Inhibitory and Antibacterial Peptides from Lactobacillus helveticus PR4 Proteinase-Hydrolyzed Caseins. *Applied and Environmental Microbiology*, 69(9), 5297-5305.
- Moatsou, G., & Sakkas, L. (2019). Sheep milk components: Focus on nutritional advantages and biofunctional potential. *Small Ruminant Research*, 180, 86-99.  
<https://doi.org/10.1016/j.smallrumres.2019.07.009>
- Moises, R. M M., Salgueiro, C. C. M., Leitão, M. S. P., & Nunes, J. F. (2024). Exploring goat's milk cheese: A systematic review of production techniques and innovations (2013-2023). *Brazilian Journal of Food Technology*, 27, e2023152.  
<https://doi.org/10.1590/1981-6723.15223>
- Morr, C.V., & Ha, E.Y.W. (1993). Whey protein concentrates and isolates: processing and functional properties. *Critical Reviews in Food Science and Nutrition*, 33(6), 431-431.
- Muguerza, B., Ramos, M., Sánchez, E., Manso, M. A., Miguel, M., Aleixandre, A., Recio, I. (2006). Antihypertensive activity of milk

- fermented by *Enterococcus faecalis* strains isolated from raw milk. *International Dairy Journal*, 16(1), 61-69.
- Nagpal, R., Kumar, A., Kumar, M., Behare, P. V., Jain, S., & Yadav, H. (2012). Probiotics, their health benefits and applications for developing healthier foods: a review. *FEMS microbiology letters*, 334(1), 1-15.
- Olveira, G., & González-Molero, I. (2016). Actualización de probióticos, prebióticos y simbióticos en nutrición clínica. *Endocrinología y Nutrición*, 63(9), 482-494.
- Parra, R. A. (2012). Yogur en la salud humana. *Revista lasallista de investigación*, 9, 162-177.
- Park, Y.W. (2006). Goat milk. Chemistry and nutrition. En Y.W. Park & G.F.W. Haenlein, *Handbook of milk of non-bovine mammals*, 34-58.
- Park, Y. W., Juárez, M., Ramos, M., & Haenlein, G. F. W. (2007). Physico-chemical characteristics of goat and sheep milk. *Small Ruminant Research*, 68(1-2), 88-113.
- Pérez, D. y Sánchez, L. (2012). Leches fermentadas: Aspectos nutritivos, tecnológicos y probióticos más relevantes. Agencia Aragonesa de Seguridad Alimentaria. Available at [https://www.aragon.es/documents/20127/674325/LECHES\\_FERMENTADAS.pdf/62a7dc81-fd9d-a459-c789-b312b658f459](https://www.aragon.es/documents/20127/674325/LECHES_FERMENTADAS.pdf/62a7dc81-fd9d-a459-c789-b312b658f459)
- Pintado, V. & Parra, J. (2012). *Elaboración de manjar utilizando suero de quesería a diferentes niveles como sustituto de la leche en el cantón Pastaza*. [Tesis de grado]. Universidad Estatal Amazónica.
- Plata, A., Ramírez, S., & Riaño Luna, C. E. (2013). Valor agregado para el lacto suero de caprino proveniente de agroindustrias lácteas: ácido láctico como alternativa de descontaminación. *REVISTA NOVA* , 11(19). <https://doi.org/10.22490/24629448.1023>

- Pordesimo, L.O. y Onwulata, C.I. (2008). *Whey texturization for snacks*. En C.I. Onwulata & P.J. Huth, *Whey processing, functionality and health benefits* (169-184). Iowa: Wiley-Blackwell IFT press.
- Prados, A., Gómez, S., Nova, E., & Marcos, A. (2015). El papel de los probióticos en el manejo de la obesidad. *Nutrición Hospitalaria*, 31(1), 10-18.
- Puertollano, E., Puertollano, M. A, Cruz-Chamorro, L., Alvarez de Cienfuegos, G., Ruiz-Bravo, A., y de Pablo, M. (2008). Orally administered *Lactobacillus plantarum* reduces pro-inflammatory interleukin secretion in sera from *Listeria monocytogenes* infected mice. *The British Journal of Nutrition*, 99(4), 819-825.
- Riera, F., Alvarez, A., Muñoz, D. (2004). Avances en el fraccionamiento de proteínas del lactosuero no desnaturalizadas. *Alimentacion: equipos y tecnología*, 23(192), 47-56.
- Ricci-Cabello, I., Herrera, M. O., & Artacho, R. (2012). Possible role of milk-derived bioactive peptides in the treatment and prevention of metabolic syndrome. *Nutrition Reviews*, 70(4), 241-255.
- Robin, O., Turgeon, S., & Paquin, P. (1993). Functional properties of milk proteins. En Y.H. Hui, *Dairy science and technology handbook*, 1, 277-353.
- Robinson P. K. (2015). Enzymes: principles and biotechnological applications. *Essays in biochemistry*, 59, 1-41.  
<https://doi.org/10.1042/bse0590001>
- Sampablo, V. (2019). *Leches fermentadas, tradición e innovación*. [Tesis de grado]. Universidad Complutense de Madrid
- Sánchez, B., De los Reyes-Gavilán, C., Margolles, A., & Gueimonde, M. (2009). Probiotic fermented milks: present and future. *Int J Dairy Technol*, 62, 472-483.

- Sanmartín, B. (2010). *Aprovechamiento de suero de quesería de origen caprino mediante la obtención de concentrados de proteínas séricas y subproductos de clarificación. estudio de sus propiedades tecnológicas.* . [Tesis de grado]. Universidad de Santiago de Compostela
- Santillán-Urquiza, E., Méndez-Rojas, M. A., & Ruiz, J. V. (2014). Productos lácteos funcionales, fortificados y sus beneficios en la salud humana. *Temas Selectos De Ingeniera En Alimentos*, 8(1), 5-14.
- Scott, R. (1991). *Fabricación de queso*. Zaragoza: Acribia S.A.
- Tirado, D. F., Acevedo, D., y Montero P. M. (2015). Extracción de Proteínas del Lactosuero de la Leche de Cabra Mediante la Aplicación de Campos Eléctricos Pulsantes de Alta Intensidad (CEPAI) . *Información Tecnológica*, 26(5).  
<https://doi.org/10.4067/S0718-07642015000500010>
- Tunick, M.H. (2008). *Whey protein production and utilization: a brief history*. En C.I. Onwulata & P.J. Huth, *Whey processing, functionality and health benefits* (1-13). Iowa: Wiley-Blackwell IFT press.
- Ulsenheimer, B. C., Pereira, D. I. B., Costa, M. M. da ., Santos, H. F. do., Peixoto, R. de M., Sangioni, L. A., & Botton, S. de A. (2022). Biosecurity in dairy goat farms. *Ciência Rural*, 52(8), e20210315.  
<https://doi.org/10.1590/0103-8478cr20210315>
- U.S. Department of Health and Human Services. (2019). *Grade "A" Pasteurized Milk Ordinance*. [https://www.ams.usda.gov/sites/default/files/media/FM MO IDFA 340.pdf](https://www.ams.usda.gov/sites/default/files/media/FM%20IDFA%20340.pdf)
- Valencia, J. (2009). El Suero de Quesería y sus Posibles Aplicaciones. *Mundo Lácteo y Cárnico.Subsecretaria de Agroindustrias y Mercados* , (2),4-6
- Verhel, A. L. (2015). *Elaboración de leche condensada de leche de bufalla (Bubalus bubalis) adicionada con oligofructosa*. [Tesis de Maestría]. Universidad Nacional de Colombia

- Vilaplana, M. (2015). Probióticos y salud. *Farmacia Profesional* 29(3), 36-39.
- Villanueva-Flores, R. (2015). Probióticos: una alternativa para la industria de alimentos. *Ingeniería Industrial*, (33), 265-275.
- Walstra, P., Wouters, J.T.M., & Geurts, T.J. (2006). *Dairy science and technology*. United States of America: CRC Press Taylor & Francis Group.
- Walzem, R.L., Dillard, C.J., & German, J.B. (2002). Whey components: Millennia of evolution create functionalities for mammalian nutrition: What we know and what we may be overlooking. *Critical Reviews in Food Science and Nutrition*, 42(4), 353-375.
- Walzem, R. (2004). *Nutritional properties of whey, lactose, and milk minerals products: Nutritional properties of whey products*. En J. Page, D. Meyer, B. Haines, V. Lagrange, & A. Kenney, *Reference manual for U.S. whey and lactose products*, 53-57.
- Webb, A. (2024). Applications of fractional stochastic volatility models to market microstructure theory and optimal execution strategies. *Frontiers in Applied Mathematics and Statistics*, 10, e 1456746. <https://doi.org/10.3389/fams.2024.1456746>
- Zehren, V. (1976). *Manual de Tecnología quesera*. Laboratorio de Tecnología del Uruguay. Montevideo.
- Zhou, G., & Sun, J. (2010). "Study on Effective Microorganisms Bacteria for Acrylonitrile Wastewater Treatment," 2010 4th International Conference on Bioinformatics and Biomedical Engineering, Chengdu, China, 2010, 1-4. <https://doi.org/10.1109/ICBBE.2010.5517301>



This edition of "*Chemical and food processes applied to goat's milk: Non-Conventional Derivatives*" was completed in the city of Colonia del Sacramento in the Eastern Republic of Uruguay on February 20, 2025



# CHEMICAL AND FOOD PROCESSES APPLIED TO GOAT'S MILK: NON-CONVENTIONAL DERIVATIVES

ISBN: 978-9915-9752-9-0



9 789915 975290